Adult Langerhans cell histiocytosis: a rare cause of colon polyp

Fig. 1 Endoscopic view of the Langerhans cell histiocytosis polyp.

A 65-year-old man with no prior medical problems underwent a screening colonoscopy. He was an ex-smoker with a 100 pack-year cigarette smoking history. A 6-mm sessile, benign-appearing polyp was removed from the descending colon using snare diathermy (● **Fig. 1**).

No other lesions were noted on colonoscopy. Histology showed a nodular proliferation of histiocytic cells within the submucosa, accompanied by a moderate number of eosinophils and lymphoid cells (● **Fig. 2** and **3**).

Immunoperoxidase studies showed that the histiocytic cells were reactive to S100, CD68, and CD1a antigens (● **Fig. 4**). Based on these results, a diagnosis of Langerhans cell histiocytosis was made. Chest and skull radiographs, bone scan, and an abdominal ultrasound were unremarkable. High-resolution computed tomography of the chest, and bone marrow aspiration and biopsy were normal. As the patient was clinically asymptomatic and the entire work-up did not reveal any other organ involvement, no therapy was initiated; 1 year later, he remains asymptomatic on regular follow-up.

Langerhans cell histiocytosis is rare in adults [1], and gastrointestinal tract involvement is even rarer [2–4]. With the increasing number of colonoscopies being performed, gastroenterologists should be aware of this rare cause of colon polyps to ensure proper follow-up and further testing to rule out systemic disease.

Endoscopy_UCTN_Code_CCL_1AD_2AC

Fig. 2 Medium magnification image of benign colonic mucosa overlying core of polyp, exhibiting pale histiocytic cells and eosinophils (hematoxylin-phloxine-safranin stain, $\times 100$).

Fig. 3 High-magnification image of colonic polyp, exhibiting histiocytic cells with pale cytoplasm and admixed eosinophils (hematoxylin-phloxine-safranin stain, $\times 400$).

Fig. 4 Immunoperoxidase reaction of lesional cells with anti-CD1a antibodies, characteristic of Langerhans cells (anti-CD1a with hematoxylin counterstain, $\times 400$).

R. Kibria¹, P. M. Gibbs², D. M. Novick¹

¹ Gastroenterology, Wright State University School of Medicine, Dayton, Ohio, USA

² Anatomical and Clinical Pathology, Wright State University School of Medicine, Dayton, Ohio, USA

References

- 1 Stocksclaeder M, Sucker C. Adult Langerhans cell histiocytosis. *Eur J Haematol* 2006; 76: 363–368
- 2 Sharma S, Gupta M. A colonic polyp due to Langerhans cell histiocytosis: a lesion not to be confused with metastatic malignant melanoma. *Histopathology* 2006; 49: 438–439
- 3 Nihei K, Terashima K, Aoyama K et al. Benign histiocytosis X of stomach. Previously undescribed lesion. *Acta Pathol Jpn* 1983; 33: 577–588
- 4 Iwafuchi M, Watanabe H, Shiratsuka M. Primary benign histiocytosis X of the stomach. A report of a case showing spontaneous remission after 5 1/2 years. *Am J Surg Pathol* 1990; 14: 489–496

Bibliography

DOI 10.1055/s-0029-1214689

Endoscopy 2009; 41: E160–E161

© Georg Thieme Verlag KG Stuttgart · New York ·

ISSN 0013-726X

Corresponding author

R. Kibria, MD

Gastroenterology

Wright State University School of Medicine

4100 W. Third Street

Dayton

Ohio 45428

USA

Fax: +1-937-268-6511

rekibria@gmail.com