Treatment of Dieulafoy's lesion of the right colon with epinephrine injection and argon plasma coagulation

Dieulafoy's lesion is a tiny submucosal defect overlying an artery in the muscularis mucosa [1]. Dieulafoy's lesion of the colon is a rare cause of lower gastrointestinal bleeding [2].

A 63-year-old woman with multiple myeloma underwent autologous bone marrow transplantation and after 6 weeks developed massive hematochezia with hemodynamic instability. Colonoscopy demonstrated bright red blood in the terminal ileum, all of the colon, and the rectum (● Fig. 1).

After the area had been washed with water, a point of spurting active bleeding was located in the ascending colon. We injected epinephrine and the bleeding stopped; identification of a minute mucosal defect was then possible (● Fig. 2).

We complemented the treatment with argon plasma coagulation (APC) using a 2.3-mm probe, with flow rate of 1.0 L/minute and a setting of 40 W in order to minimize the risk of bowel perforation, until the lesion was completely coagulated. Submucosal injection of epinephrine has a protective effect when using thermal techniques in the right colon [3]. There was no rebleeding during the follow-up of 14 days. The patient died after this period from septic shock.

Dieulafoy's lesion in the setting of hemorrhagic shock has a high risk of rebleeding, justifying the addition of a complementary endoscopic treatment (thermal or mechanical) following epinephrine injection [4–5]. However, we have to keep in mind that the right colon has a thinner wall compared with the stomach, and use of band ligation and heat probe with high temperatures can lead to bowel perforation.

Fig. 1 Colonoscopy showing spurting active bleeding in ascending colon.

Fig. 2 Minute mucosal defect in ascending colon identified after injection of epinephrine.

To our knowledge, this is the first report of a combined endoscopic approach with injection of epinephrine and APC to treat a Dieulafoy's lesion of the right colon in a patient with significant thrombocytopenia. It seems to be a secure and effective modality of endoscopic therapy of bleeding in this setting, with minimum risk of perforation and high possibility of hemostasis.

Endoscopy_UCTN_Code_CPL_1AH_2AB

J. L. S. Souza

Department of Gastroenterology,
Diagnostic Center in Gastroenterology,
University of Sao Paulo, Brazil

References

- 1 Juler GL, Labitzke HG, Lamb R et al. The pathogenesis of Dieulafoy's gastric erosion. *Am J Gastroenterol* 1984; 79: 195–200
- 2 Norton ID, Petersen BT, Sorbi D et al. Management and long term prognosis of Dieulafoy lesion. *Gastrointest Endosc* 1999; 50: 762–767
- 3 Suzuki N, Arebi N, Saunders BP. A novel method of treating colonic angiodysplasia. *Gastrointest Endosc* 2006; 64: 424–427
- 4 Calvet X, Vergara M, Brullet E et al. Addition of a second endoscopic treatment following epinephrine injection improves outcome in high risk bleeding ulcers. *Gastroenterology* 2004; 126: 441–450
- 5 Chung IK, Kim EJ, Lee MS et al. Bleeding Dieulafoy's lesions and the choice of endoscopic method: comparing the hemostatic efficacy of mechanical and injection methods. *Gastrointest Endosc* 2000; 52: 721–724

Bibliography

DOI 10.1055/s-0029-1214774

Endoscopy 2009; 41: E192

© Georg Thieme Verlag KG Stuttgart · New York ·
ISSN 0013-726X

Corresponding author

J. L. S. de Souza, MD

Department of Gastroenterology,
Diagnostic Center in Gastroenterology,
University of São Paulo
255 Dr Enéas de Carvalho Aguiar Ave.
9th floor – Room 9159
São Paulo
Brazil
Fax: +55-11-30697940
jlsebba@gmail.com