Anisakiasis and vanishing tumor of the cecum

Fig. 1 Colonoscopy showing a lesion in the cecum, similar to a Borrmann type 2 lesion.

Fig. 2 Histopathological section of the biopsy specimen showing eosinophilic infiltration in the lamina propria mucosae and necrosis of the epithelium.

There have been few reports to date of vanishing gastric tumors caused by anisakiasis [1]. The large intestine is rarely affected because the ingested larvae usually do not travel that far down the gastrointestinal tract. Colonic anisakiasis occasionally resembles a colonic tumor, because it leads to edema, acute phlegmonous inflammation, or the formation of granulomas around the larvae in the submucosa of the intestinal wall [2].

A 77-year-old man attended our hospital for a positive fecal occult blood test. He was asymptomatic. He had a history of appendectomy for acute appendicitis at the age of 19 and ate raw fish almost every day. His white blood cell count was normal with no eosinophilia. At colonoscopy, a clearly demarcated, depressed lesion with raised margins (similar to a Borrmann type 2 lesion) was identified in the cecum (● Fig. 1).

The histopathological examination of the biopsy specimens revealed eosinophilic infiltration of the lamina propria mucosae and the submucosa, and necrosis of the epithelium (● Fig. 2).

Malignant cells were not seen. Computed tomography showed wall thickening in the cecum (● Fig. 3).

On repeat colonoscopy after 16 days, the tumour resembling a Borrmann type 2 lesion had disappeared and a reddish scar with small erosion was seen in the cecum (● Fig. 4).

Serum titers of both IgG and IgA antibodies to *Anisakis* larvae on the day of the second colonoscopy were slightly elevated at 1.61 (cut-off index, normal < 1.50). At another colonoscopy a year later, the

Fig. 3 Computed tomography scan showing cecal wall thickening and the "dirty fat" sign around the lesion.

Fig. 4 Repeat colonoscopy showing disappearance of the tumor and a reddish scar with a small erosion in the cecum.

Fig. 5 Colonoscopy after a year showing a whitish scar but no evidence of tumor.

tumor had completely disappeared and a whitish scar was seen in the cecum (● Fig. 5).

The vanishing tumor may be considered to be anisakiasis of the cecum.

Endoscopy_UCTN_Code_CCL_1AD_2AZ

**N. Ishii, M. Matsuda, T. Setoyama,
S. Suzuki, S. Uchida, M. Uemura,
Y. Iizuka, K. Fukuda, N. Horiki, Y. Fujita**
Department of Gastroenterology,
St. Luke's International Hospital, Tokyo,
Japan

References

- 1 Takeuchi K, Hanai H, Iida T et al. A bleeding gastric ulcer on a vanishing tumor caused by anisakiasis. *Gastrointest Endosc* 2000; 52: 549–551
- 2 Verhamme MA, Ramboer CH. Anisakiasis caused by herring in vinegar: a little known medical problem. *Gut* 1988; 29: 843–847

Bibliography

DOI 10.1055/s-0029-1214923

Endoscopy 2009; 41: E226–E227

© Georg Thieme Verlag KG Stuttgart · New York ·
ISSN 0013-726X

Corresponding author

N. Ishii

Department of Gastroenterology
St. Luke's International Hospital
9-1 Akashi-cho
Chuo-ku
Tokyo
Japan 104-8560
Fax: +81-3-35440649
naoishi@luke.or.jp