Lansoprazole-associated collagenous colitis: unique presentation, similar to ischemic colitis

Fig. 1 Abdomino-pelvic computed tomography (CT) scan demonstrating the thick-walled sigmoid colon.

Fig. 3 Biopsy specimen showing markedly thickened subepithelial collagen bands.

Fig. 2 Colonoscopy showing an actively hemorrhagic linear ulcer and a linear ulcer scar.

Fig. 4 Follow-up colonoscopy revealed the complete healing of the linear ulcer.

Recently, there have been reports of lansoprazole-associated collagenous colitis manifesting some characteristic clinical findings: chronic diarrhea, typical endoscopic findings, pathological evidence of collagen bands, and rapid clinical improvement after discontinuation of lansoprazole. We have also often encountered patients taking lansoprazole presenting with chronic diarrhea and typical endoscopic findings. Although one published article has stated that linear ulcers and ulcer scars should be considered specific for lansoprazole-associated collagenous coli-

tis, another has not mentioned endoscopic findings at all [1,2]. However, severe ulcers rarely cause colonic perforation [3]. Here we present a patient with lansoprazole-associated collagenous colitis who presented a unique clinical picture, similar to that of ischemic colitis. A 78-year-old Japanese woman attended our emergency room because of abrupt onset of lower abdominal pain and heavy blood in her stool. She had been taking lansoprazole since 2 months and had chronic diarrhea. Initially, ischemic colitis was suspected and an abdominopelvic

computed tomography (CT) scan revealed a thick-walled and edematous sigmoid colon (● Fig. 1). Colonoscopy was performed to confirm the diagnosis and revealed a 20-cm long hemorrhagic, linear ulcer and a 15-cm linear ulcer scar in the sigmoid colon (● Fig. 2). Histopathological examination of biopsy samples taken from the sigmoid colon showed subepithelial collagen bands (● Fig. 3). Finally, a diagnosis of collagenous colitis was made. The cause was thought to be lansoprazole and this was discontinued. Six days later, the patient was discharged with complete resolution of the diarrhea and abdominal pain. A repeat colonoscopy 2 months after admission showed healing linear lesions and scars (● Fig. 4). No collagen bands were identified on a biopsy specimen.

The present case highlights the fact that lansoprazole-associated collagenous colitis may present with atypical clinical, imaging, and endoscopic findings.

Endoscopy_UCTN_Code_CCL_1AD_2AJ

H. Yusuke, T. Jun, M. Naotaka, T. Yuichi, E. Yutaka, I. Kazuaki

Department of Internal Medicine,
Fujigaoka Hospital, Showa University,
Yokohama, Japan

References

- 1 Umeno J, Matsumoto T, Nakamura S et al. Linear mucosal defect may be characteristic of lansoprazole-associated collagenous colitis. *Gastrointest Endosc* 2008; 67: 1185–1191
- 2 Thomson RD, Lestina LS, Bensen SP et al. Lansoprazole-associated microscopic colitis: a case series. *Am J Gastroenterol* 2002; 97: 2908–2913
- 3 Smith RR, Ragput A. Mucosal tears on endoscopic insufflation resulting in perforation: an interesting presentation of collagenous colitis. *J Am Coll Surg* 2007; 205: 725

Bibliography

DOI 10.1055/s-0029-1214942

Endoscopy 2009; 41: E281–E282

© Georg Thieme Verlag KG Stuttgart · New York ·
ISSN 0013-726X

Corresponding author

Y. Hashimoto, MD

Department of Internal Medicine

Fujigaoka Hospital

Showa University

1-30 Aoba-ku Fujigaoka

Yokohama

Japan 227-8501

Fax: +81-045-9713824

yu-hashimoto@showa-university-fujigaoka.gr.jp