A blue patient with patent blue

A 68-year-old man underwent colonoscopy because of a 3-cm semicircular sessile polyp in the transverse colon, close to the left flexure. The aim was to mark the polyp for the operative resection planned for the next day. To mark the polyp, 6 ml patent blue was used (Patentblau V; Guerbet GmbH, Sulzbach, Germany). As a first step, depots of saline solution were injected at the edges of the polyp. After that, the patent blue solution was injected.

Fifteen minutes after the end of the colonoscopy, the nurse observed that the patient's lips were blue, without any sign of dyspnea. For safety reasons he was not removed to the ward for another 3 hours. In these 3 hours the color of his skin changed to dark blue all over his body (► Fig. 1). By 48 hours after the injection, the dark blue color had vanished and the skin was pink again. During the whole time the patient was very well, without any signs of dyspnea or allergic reaction.

Patent blue is a substance which is increasingly being used to mark the sentinel node and for lymphatic mapping in carcinoma of the colon and rectum [1,2]. It is also used as a food colorant (e.g., blue curacao, jelly beans). There have been a few cases of localized adverse events occurring during mapping for breast cancer and melanoma, but no discoloring of the whole body and none at all when patent blue was being used in the colon [3]. Two reasons are suspected as the cause of the blue skin color: discoloration due to lymphatic spread from the injection site, and methemoglobinemia [4]. Anaphylactic reactions have been described as a side effect, and intracutaneous tests with the dye should be done in advance of any endoscopic procedure in which the dye is used [5].

Endoscopy_UCTN_Code_CPL_1AJ_2AI

Fig. 1 Blue skin color over a patient's whole body 3.25 hours after injection of patent blue, performed to mark a colonic polyp intended for resection the following day

S. Grüne, E. Treher

Department of Gastroenterology, Hospital Hetzelstift, Neustadt an der Weinstraße, Germany

Bibliography

DOI 10.1055/s-0029-1215412

Endoscopy 2010; 42: E11

© Georg Thieme Verlag KG Stuttgart · New York · ISSN 0013-726X

References

- 1 Cahill RA, Leroy J, Marescaux J. Could lymphatic mapping and sentinel node biopsy provide oncological providence for local resectional techniques for colon cancer? *BMC Surg* 2008; 8: 17
- 2 Kelder W, Braat AE, Karrenbeld A et al. The sentinel node procedure in colon carcinoma: a multi-centre study in The Netherlands. *Int J Colorectal Dis* 2007; 22: 1509–1514
- 3 Daley MD, Norman PH, Leak JA et al. Adverse events associated with the intraoperative injection of isosulfan blue. *J Clin Anesth* 2004; 16: 332–341
- 4 Yusim Y, Livingstone D, Sidi A. Blue dyes, blue people: the systemic effects of blue dyes when administered via different routes. *J Clin Anesth* 2007; 19: 315–321
- 5 Scherer K, Studer W, Figueiredo V, Bircher AJ. Anaphylaxis to isosulfan blue and cross-reactivity to patent blue V: case report and review of the nomenclature of vital blue dyes. *Ann Allergy Asthma Immunol* 2006; 96: 497–500

Corresponding author

S. Grüne, MD

Department of Gastroenterology
Hospital Hetzelstift
Stiftstraße 10
67434 Neustadt an der Weinstraße
Germany
Fax: +49-6321-8594007
Grüne@ekgsg.de