Esophagomediastinal and esophagobronchial fistulas associated with invasive aspergillosis

A 67-year-old woman was diagnosed as having esophageal cancer. She was given neoadjuvant chemoradiotherapy and esophagectomy was carried out. Reconstruction was accomplished with a gastric pull-up through the posterior mediastinal route. After 6 years of the treatment, the patient had high grade fever, which was refractory to treatment with antibiotics. A chest computed tomography (CT) scan demonstrated mixed fluid and soft tissue density in the posterior mediastinum, partially within the wall of the esophagus (● Fig. 1). A Gastrografin esophagogram confirmed the existence of esophagomediastinal and esophagobronchial fistulas (● Fig. 2). We considered these fistulas as the cause of the fever, and carried out percutaneous drainage. Gastrointestinal endoscopy revealed an esophagomediastinal fistula with necrotic tissues (● Fig. 3). Although both sputum and blood cultures did not reveal the causative organism, high levels of serum galactomannan (whose production is proportional to the *Aspergillus* fungal load in tissue) were demonstrated (> 5.0 ng/mL; cut-off value < 0.5 ng/mL). On the basis of this finding, along with the persistent fever and CT appearances, a diagnosis of probable invasive aspergillosis was made in accordance with the European Organisation for the Research and Treatment of Cancer/Mycoses Study Group (EORTC/MSG) consensus criteria [1]. Treatment with antifungal agents led to a fall in the temperature, reduction in the hematological parameters, including galactomannan, and resolution of the radiological findings. At 1 month, endoscopy revealed healing of the fistula (● Fig. 4).

Aspergillus is a common airborne organism that can be highly pathogenic under immunocompromised conditions such as prolonged neutropenia after chemotherapy or organ transplantation [2]. Mediastinitis caused by *Aspergillus* infection usually occurs by airborne contamination of the cardiothoracic surgical field in the months following the surgery [3]. Invasive aspergillosis is a fairly rare condition, but

Fig. 1 A fluid and soft tissue density seen in the posterior mediastinum, which was not separable from the reconstructed esophagus.

Fig. 2 Gastrografin esophagogram showing perforation of the reconstructed esophagus with contrast flowing into the mediastinum and into the bronchial tree in the upper lobe of the right lung.

Fig. 3 Gastrointestinal endoscopy showing an esophagomediastinal fistula surrounded by necrotic tissue. A drainage tube was inserted into the fistula via the percutaneous route.

Fig. 4 At 1 month after treatment with antifungal agents and percutaneous drainage, endoscopy showed healing of the fistula.

can be devastating if there is a delay in diagnosis [4]. This diagnosis needs to be considered in a patient with antibiotic-refractory infection in the presence of local immunocompromising factors such as irradiation-induced lung injury.

Endoscopy_UCTN_Code_CCL_1AB_2AC_3AZ

S. Kato¹, S. Inoue², M. Inamori¹,
N. Miyazawa², T. Sato², N. Kobayashi²,
K. Mishina², M. Sasaki², Y. Ishigatsubo²,
A. Nakajima¹

¹ Gastroenterology Division, Yokohama City University, School of Medicine, Japan

² Department of Internal Medicine and Clinical Immunology, Yokohama City University, School of Medicine, Japan

References

- 1 Ascioglu S, Rex JH, de Pauw B et al. Defining opportunistic invasive fungal infections in immunocompromised patients with cancer and hematopoietic stem cell transplants: an international consensus. *Clin Infect Dis* 2002; 34: 7–14
- 2 Latge JP. *Aspergillus fumigatus* and aspergillosis. *Clin Microbiol Rev* 1999; 12: 310–350
- 3 Levin T, Suh B, Beltramo D et al. *Aspergillus* mediastinitis following orthotopic heart transplantation: case report and review of the literature. *Trans Infect Dis* 2004; 6: 129–131
- 4 Hope WW, Waish TJ, Denning DW. Laboratory diagnosis of invasive aspergillosis. *Lancet Infect Dis* 2005; 5: 609–622

Bibliography

DOI 10.1055/s-0029-1215414

Endoscopy 2010; 42: E48–E49

© Georg Thieme Verlag KG Stuttgart · New York · ISSN 0013-726X

Corresponding author

S. Kato, MD

Gastroenterology Division
Yokohama City University Hospital
3-9 Fukuura
Kanazawa-ku
Yokohama 236-0004
Japan
Fax: +81-45-7843546
shin800m@yokohama-cu.ac.jp