S. SHINOZAKI, T. HAMURA, Y. IBUSUKI, K. FUJII, H. UEKUSA, K. SUZUKI* (SORST, JST AGENCY, TOKYO, TOKYO INSTITUTE OF TECHNOLOGY, KWANSEI GAKUIN UNIVERSITY, SANDA AND PRESTO, JST AGENCY, TOKYO, JAPAN)
 Hexaradialenes by Successive Ring Openings of Tris(alkoxy)tricyclobutabenzenes: Synthesis and Characterization
Angew. Chem. Int. Ed. **2010**, *49*, 3026–3029.

Radialene Synthesis

Significance: The authors present a remarkable synthetic route to hexaradialene **1** that utilizes a thermally-driven ring-opening isomerization reaction to form the six exocyclic double bonds in the final step. The reaction proceeds efficiently provided that sufficiently bulky groups are tethered to the triannulated parent compound prior to isomerization, with both the *syn* and the *anti* precursors converted into **1**.

Comment: Alicyclic cross-conjugated molecules of this type are rare and further studies on the reactivity of these intriguing molecules could provide new routes to novel polycyclic aromatic systems.

SYNFACTS Contributors: Timothy M. Swager, D. Barney Walker
 Synfacts 2010, 6, 0649-0649 Published online: 20.05.2010
 DOI: 10.1055/s-0029-1219982; Reg-No.: S04910SF