Fatal venous air embolism during upper endoscopy in a patient with percutaneous transhepatic cholangiography (PTC) catheter

Venous air embolism is a rare complication of endoscopic procedures reported with colonoscopy, esophagogastroduodenoscopy, endoscopic ultrasound with fine needle aspiration and endoscopic retrograde cholangiography with sphincterotomy [1–5]. We report a case of a fatal venous air embolism during upper endoscopy in a patient with prior pancreaticoduodenectomy and recent percutaneous transhepatic cholangiography (PTC) catheter placement.

A 52-year-old man underwent pancreaticoduodenectomy for pancreatic adenocarcinoma and PTC catheter placement for ascending cholangitis. An upper endoscopy under fluoroscopy was performed after the patient developed melena. The endoscope was advanced to the hepaticojejunostomy (● Fig. 1), when the patient suddenly became hypoxic, bradycardic, and developed pulseless electrical activity. Despite prolonged cardiopulmonary resuscitation, the patient expired. An autopsy was declined by the family, however, review of the intraoperative fluoroscopic images revealed an air venogram, consistent with a fatal venous air embolism (● Fig. 2).

Occurrence of venous air embolism during endoscopy requires communication between the air source and vasculature as well as a pressure gradient favoring passage of air into the venous circulation. Air usually enters through a mucosal defect or vasculoenteric fistula, aided by positive pressure from insufflation. Clinical manifestations include sudden decrease in end-tidal carbon dioxide or oxygen saturation, cyanosis, hypotension, bradycardia or tachycardia, and a mill-wheel murmur. Initial management consists of provision of high-flow oxygen, Trendelenburg position, and insertion of a central venous catheter for aspiration of gas bubbles. Although venous air embolism may occur in patients without risk factors [6, 7], disruption of mucosal barriers (ulcerations, trauma, dilation, sphincterotomy), presence of blind intestinal limbs (hepaticojejunostomy), and bowel directly apposed to venous structures (varices) are known risk factors [8–12]. In the present patient, placement of the PTC catheter probably

created a vasculobiliary fistula. The presence of a widely patent hepaticojejunostomy, blind intestinal limb, and air insufflation created a pressure gradient allowing air flow into the venous circulation. The fluoroscopic images confirmed air within the hepatic veins and inferior vena cava (● Fig. 2). General endotracheal anesthesia and close monitoring of endotracheal carbon dioxide in patients with known risk factors may improve clinical outcomes.

Endoscopy_UCTN_Code_CPL_1AH_2AB
Endoscopy_UCTN_Code_CPL_1AK_2AH

C. B. Meier¹, A. J. Moser², M. K. Sanders³

¹ University of Colorado Health Sciences Center – Gastroenterology, Hepatology, Aurora, Colorado, USA

² Department of Surgery, University of Pittsburgh Medical Center, Pittsburgh, Pennsylvania, USA

³ Division of Gastroenterology, Hepatology, and Nutrition, University of Pittsburgh Medical Center, Pittsburgh, Pennsylvania, USA

References

- Huycke A, Moeller DD. Hepatic portal venous gas after colonoscopy in granulomatous colitis. *Am J Gastroenterol* 1985; 80: 637–638
- Akhtar N, Jafri W, Mozaffar T. Cerebral artery air embolism following an esophagogastros-copy: a case report. *Neurology* 2001; 56: 136–137
- Patel RT, Deen KI, Fielding JW. Portal pyaemia following flexible sigmoidoscopy. *Br J Surg* 1995; 82: 422–423
- Pfaffenbach B, Wegener M, Bohmeke T. Hepatic portal venous gas after transgastric EUS-guided fine-needle aspiration of an accessory spleen. *Gastrointest Endosc* 1996; 43: 515–518
- Kennedy C, Larvin M, Linsell J. Fatal hepatic air embolism following ERCP. *Gastrointest Endosc* 1997; 45: 187–188
- Demaerel P, Gevers AM, De Bruecker Y et al. Stroke caused by cerebral air embolism during endoscopy. *Gastrointest Endosc* 2003; 57: 134–135
- Chorost MI, Wu JT, Webb H, Ghosh BC. Vertebral venous air embolism: an unusual complication following colonoscopy: report of a case. *Dis Colon Rectum* 2003; 46: 1138–1140
- Huycke A, Moeller DD. Hepatic portal venous gas after colonoscopy in granulomatous colitis. *Am J Gastroenterol* 1985; 80: 637–638

Fig. 1 Endoscope advanced through the afferent limb to the hepaticojejunostomy.

Fig. 2 Venous air embolism.

- Katzgraber F, Glenewinkel F, Rittner C, Beule J. Fatal air embolism from gastroscopy [letter]. *Lancet* 1995; 346: 1714–1715
- Barthet M, Membrini P, Bernard J-P, Sahel J. Hepatic portal venous gas after endoscopic biliary sphincterotomy. *Gastrointest Endosc* 1994; 40: 261–263
- Lowdon Jd, Tidmore TL. Fatal air embolism after gastrointestinal endoscopy. *Anesthesiology* 1988; 69: 622–623
- Desmond PV, MacMahon RA. Fatal air embolism following endoscopy of a hepatic portoenterostomy [letter]. *Endoscopy* 1990; 22: 236

Bibliography

DOI 10.1055/s-0029-1243942

Endoscopy 2010; 42: E111

© Georg Thieme Verlag KG Stuttgart · New York · ISSN 0013-726X

Corresponding author

C. B. Meier

University of Colorado Health Sciences Center – Gastroenterology, Hepatology
12631 East 17th Avenue
MS B158 Aurora
Colorado 80045
USA
+1-720-848-2756
meiercb@hotmail.com