Small-bowel obstruction precipitated by bowel preparation for screening colonoscopy

A 55-year-old man with a remote history of appendectomy presented to the emergency department 24 hours after undergoing a screening colonoscopy, complaining of persistent nausea, emesis, and worsening abdominal pain that began during a split-day bowel preparation with low-volume polyethylene glycol and bisacodyl tablets. The patient reported that he developed nausea and generalized abdominal cramping after he finished the first part of bowel preparation and then developed multiple episodes of emesis after the second part. However, his vital signs and abdominal exam were unremarkable and colonoscopy was performed without difficulty. At home he had recurrent nausea, emesis, and worsening abdominal pain, and therefore went to the emergency department. His abdomen was distended, tympanitic, and diffusely tender to palpation, with hypoactive, high-pitched bowel sounds and no peritoneal signs. Plain radiographs of the abdomen revealed no free air, but did indicate distension of the proximal and mid small bowel with multiple air-fluid levels (● Fig. 1).

After conservative management failed, the patient underwent a laparotomy which revealed an obstruction in the ileum secondary to a stricture caused by an adhesion.

The most common cause of small-bowel obstruction (SBO) is adhesions (65%–75% of cases) [1]. The literature contains a few case reports of SBO secondary to colonoscopy [2–5]. Ours is the first report of a SBO induced by bowel preparation. Review of the cases of colonoscopy-induced

Fig. 1 Plain abdominal radiograph (upright) demonstrating distension of proximal and mid small bowel with multiple air-fluid levels.

SBO, as well as of this present case, found that a common factor was prior abdominal surgery. The fact that symptoms of SBO started almost immediately after the patient's bowel preparation was finished and before his arrival for the colonoscopy makes it less likely that the procedure itself induced SBO. The rare complication of SBO should be considered in a patient presenting with symptoms of nausea, vomiting, and abdominal pain at the time of colonoscopy.

Competing interests: None

Endoscopy_UCTN_Code_CPL_1AJ_2AB

J. T. Hlivko¹, E. J. Esber¹, J. A. Porter², C. H. Kefalas¹

¹ Department of Medicine, Summa Health System, Akron, Ohio, USA

² Department of Surgery, Summa Health System, Akron, Ohio, USA

References

- 1 Menzies D, Ellis H. Intestinal obstruction from adhesions – how big is the problem? *Ann R Coll Surg Engl* 1990; 72: 60–63
- 2 Zanati SA, Fu A, Kortan P. Routine colonoscopy complicated by small-bowel obstruction. *Gastrointest Endosc* 2005; 61: 781–783
- 3 Malki SA, Bassett ML, Pavli P. Small bowel obstruction caused by colonoscopy. *Gastrointest Endosc* 2001; 53: 120–121
- 4 Gonzalez RA, Avila S, Lopez-Roses L et al. Small bowel obstruction and perforation after colonoscopy. *Endoscopy* 2003; 35: 192
- 5 Raghavendran K, Novak JM, Amodeo JL et al. Mechanical small bowel obstruction precipitated by colonoscopy. *Surg Endosc* 2003; 17: 1496

Bibliography

DOI 10.1055/s-0030-1255644

Endoscopy 2010; 42: E220

© Georg Thieme Verlag KG Stuttgart · New York · ISSN 0013-726X

Corresponding author

J. T. Hlivko, MD

Department of Medicine
Summa Health System
55 Arch Street, Suite 1A
Akron
Ohio 44304
USA
Fax: +1-330-375-3760
jonhlivko@yahoo.com