Treatment of a completely obstructed colonic anastomotic stricture using a CT-guided endoscopic rendezvous technique

Fig. 1 The first attempt at endoscopic dilatation failed because it was not certain that the two endoscopes were facing each other in this rendezvous procedure.

Fig. 2 Endoscopic view of the stenotic anastomosis from below.

Fig. 3 CT-guided fluoroscopy shows the two endoscopes exactly facing each other.

Fig. 4 Balloon dilatation following successful recanalization of the stenosis.

Fig. 5 Barium enema before the planned ileostomy takedown shows a residual stricture with good passage.

Although anastomotic strictures are a recognized complication of colorectal surgery, complete colonic anastomotic obstruction from benign disease is rare and there are few reports of endoscopic management [1–4]. Here, we describe a case of successful endoscopic treatment using a novel combined antegrade-retrograde endoscopic rendezvous technique facilitated by CT-guided fluoroscopy.

A 73-year-old man presented to us with complete anastomotic obstruction following deep anterior resection for UICC stage III rectal carcinoma complicated by an anastomotic leak that had been treated by endoscopic sponge placement in the pararectal cavity [5]. We attempted to employ a rendezvous method to perforate the stenosis as described previously [1, 3, 4], with a colonoscope passed through the existing

loop ileostomy and an upper endoscope advanced to the stenosis in a retrograde fashion. The scopes, under fluoroscopy, were seen to be in proximity to each other, but a “kissing position” could not be attained (● Fig. 1).

This was because of the J-shaped configuration of the anastomosis with two dead ends on its lower aspect, neither of which could be positively identified as the anastomotic stenosis in question (● Fig. 2).

Administration of barium through both endoscopes, in addition, revealed a nonlinear alignment of the colon and showed that the oral and aboral ends were separated only by a fibrous membrane (● Fig. 1). To demonstrate the exact position of the endoscopes in a three-dimensional fashion we repeated the procedure with CT guidance. CT fluoroscopy confirmed that the tips were facing each other when the lower endoscope was placed in one of the two dead ends (● Fig. 3).

Under CT-fluoroscopic and transillumination guidance, the fibrous septum was penetrated using a biopsy forceps passed through the accessory channel of the retrograde endoscope. After visualization of the forceps by the antegrade colonoscope, a wire-guided “through-the-scope” balloon was placed at the site of the stricture and used to dilate it sequentially to 12 mm (● Fig. 4).

Within 1 month following this first intervention, the stenosis was dilated up to 20 mm in a series of five endoscopic sessions. Clinically, the patient was well with normal defecation. Barium enema confirmed good passage (● Fig. 5) and the ileostomy was closed 6 weeks after the CT-guided endoscopic intervention.

We believe that CT fluoroscopy guidance adds a measure of safety by allowing the endoscopist to visualize the stenosis and the position of the endoscopes three-dimensionally in complex situations where unambiguous identification of the stenosis is not possible in conventional fluoroscopy. Limitations of the technique include the need for either instrument exchange or a second endoscopy unit and a pre-existing ostomy to perform the rendezvous procedure.

Endoscopy_UCTN_Code_TTT_1AQ_2AF

Competing interests: None

**M. Albertsmeier¹, P. Rittler¹,
R.-T. Hoffmann², F. Spelsberg¹**

¹ Department of Surgery, University of Munich – Campus Grosshadern, Ludwig Maximilian University of Munich, Germany

² Institute of Clinical Radiology, University of Munich – Campus Grosshadern, Ludwig Maximilian University of Munich, Germany

References

- 1 Kaushik N, Rubin J, McGrath K. Treatment of benign complete colonic anastomotic obstruction by using an endoscopic rendezvous technique. *Gastrointest Endosc* 2006; 63: 727–730
- 2 De Lusong MA, Shah JN, Soetikno R, Binmoeller KF. Treatment of a completely obstructed colonic anastomotic stricture by using a prototype forward-array echoendoscope and facilitated by SpyGlass (with videos). *Gastrointest Endosc* 2008; 68: 988–992
- 3 Davies M, Satyadas T, Akle CA, Kirkham JS. Combined endoscopic approach for the management of a difficult recto-sigmoid anastomotic stricture. *Int Surg* 2004; 89: 76–79
- 4 Reddy RA, Venkatasubramaniam AK, Khurshed A et al. Dual interventional approach of endoscopic reboring in completely stenosed rectal anastomosis using radiology guidance: a novel technique. *Colorectal Dis* 2009; 11: 49–52
- 5 Weidenhagen R, Gruetzner KU, Wiecken T et al. Endoscopic vacuum-assisted closure of anastomotic leakage following anterior resection of the rectum: a new method. *Surg Endosc* 2008; 22: 1818–1825

Bibliography

DOI 10.1055/s-0030-1255692

Endoscopy 2011; 43: E5–E6

© Georg Thieme Verlag KG Stuttgart · New York ·

ISSN 0013-726X

Corresponding author

M. Albertsmeier, MD

Klinikum der Universität München –

Standort Grosshadern

Chirurgische Klinik und Poliklinik

Marchioninistr. 15

81377 München

Germany

Fax: +49-89-70955464

Markus.Albertsmeier@med.uni-muenchen.de