Acute obstructive cholangitis caused by an enterolith in a duodenal diverticulum

Fig. 1 Computed tomographic image showing a diverticulum in the second part of the duodenum.

Fig. 2 Endoscopic examination revealed a periampullary diverticulum occupied by a yellow stone. The arrow indicates the ampulla of Vater.

Duodenal diverticula are common. While they are usually asymptomatic, several complications have been reported.

A 78-year-old man was admitted to our hospital with high-grade fever (39.6°C) and jaundice. His laboratory data showed liver dysfunction and elevations of the white blood cell count and serum C-reactive protein concentration. Abdominal ultrasonography demonstrated dilated biliary ducts. Abdominal computed tomography (CT) revealed a diverticulum in the second part of the duodenum (● Fig. 1).

We suspected obstructive cholangitis. Endoscopic retrograde cholangiography (ERC) was performed. Gastrointestinal endoscopy revealed a periampullary diverticulum occupied by a yellow stone (● Fig. 2).

ERC showed no abnormalities (● Fig. 3). Therefore, we made a diagnosis of biliary tract obstruction caused by an enterolith in the periampullary duodenal diverticulum. We performed successful endoscopic removal of the enterolith (● Fig. 4).

The extracted enterolith was oval-shaped, yellowish in color, measured approximately 3 cm in diameter, and had a hard outer rim with an irregular surface (● Fig. 5).

Separate layers within the wall were appreciated and there was no evidence of any nidus, such as a fruit pit.

Infrared absorption spectrophotometry showed that the enterolith consisted mainly of deoxycholic acid (● Fig. 6).

Duodenal diverticula are commonly encountered and are generally regarded as clinically insignificant entities. They can, however, sometimes produce serious complications, including diverticulitis, perforation, hemorrhage, biliary and/or pancreatic obstruction, partial duodenal obstruction, fistula formation with adjacent organs, diarrhea secondary to blind loop syndrome, and enterolith formation [1].

Enterolith formation is known to occur within regions of stasis, such as Meckel's diverticulum or a blind loop, or as a result of stricture due to Crohn's disease or tuberculosis. Duodenal diverticula may also represent such a region of stasis [2].

In conclusion, we performed successful endoscopic removal of an enterolith in a duodenal diverticulum. Obstructive cholangitis caused by an enterolith in a periampullary diverticulum is rare, but is an important entity for endoscopists.

Competing interests: None

Fig. 3 Endoscopic retrograde cholangiography showed no abnormalities.

Fig. 4 Successful endoscopic removal of the enterolith was performed.

Fig. 5 Enterolith removed from the duodenal diverticulum.

Fig. 6 Infrared absorption spectrophotometry showed that the enterolith consisted mainly of deoxycholic acid. T, transmittance.

Endoscopy_UCTN_Code_CCL_1AZ_2AI

T. Nonaka¹, M. Inamori², T. Kessoku¹, Y. Ogawa¹, K. Imajyo¹, S. Yanagisawa¹, T. Shiba¹, T. Sakaguchi¹, A. Nakajima², S. Maeda², K. Atsukawa¹, H. Takahashi¹

¹ Department of Gastroenterology, Hiratsuka City Hospital, Hiratsuka, Japan

² Gastroenterology Division, Yokohama City University School of Medicine, Yokohama, Japan

References

- 1 *Eeckhout G, Vanstiphout J, Van Pottelbergh I et al.* Endoscopic treatment of a perforated duodenal diverticulum. *Endoscopy* 2000; 32: 991–993
- 2 *Shocket E, Simon SA.* Small bowel obstruction due to enterolith (bezoar) formed in a duodenal diverticulum: a case report and review of the literature. *Am J Gastroenterol* 1982; 77: 621–624

Bibliography

DOI 10.1055/s-0030-1255702

Endoscopy 2010; 42: E204–E205

© Georg Thieme Verlag KG Stuttgart · New York · ISSN 0013-726X

Corresponding author

M. Inamori, MD, PhD

Gastroenterology Division
Yokohama City University School of Medicine
3-9 Fukuura
Kanazawa-ku
Yokohama
236-0004 Japan
Fax: +81-45-784-3546
inamorim@med.yokohama-cu.ac.jp