Combined colonoscopy and laparoscopy to close a colonic perforation

Fig. 1 Perforation visible on colonoscopy.

Fig. 2 Perforation site manipulated with the laparoscope.

Fig. 3 Serosal transillumination of mucosal seal with endoclips.

A 79-year-old man underwent a colonoscopy with rectal adenomatous polyp removal. As the colonoscope traversed the sigmoid colon, the operator felt a small 'pop'. An abdominal computed tomography scan revealed extraluminal air in the retroperitoneum.

Surgical consultation suggested repair of the perforation by using a combined endoscopic and laparoscopic approach. A 5-mm transmural perforation was found in the sigmoid colon (► **Fig. 1**).

Endoclips (Resolution clips; Boston Scientific, Boston, Massachusetts, USA) were applied to the mucosal side of the perforation with a colonoscope as well as laparoscopic assistance. The laparoscope was used to manipulate the colon externally from the serosal side, bringing the perforation to face the colonoscope directly, and avoiding tangential placement of the endoclips (► **Fig. 2**).

Three endoclips were deployed without complication (► **Fig. 3**).

A water irrigation test was performed: the proximal bowel was clamped and then the area of repaired perforation was placed under water from the serosal side laparoscopically.

The colonoscope was then used to insufflate the bowel with air to distend it. Considerable precautions were taken to avoid air escaping from the site of the perforation. Subsequently, small numbers of air bubbles were noted. Two sutures were

placed laparoscopically from the serosal side to seal the air leak. A second water irrigation test confirmed the closure of the perforation.

To our knowledge, this is the first case of colonic perforation where colonoscopy and laparoscopy were used to complete a repair effectively, without colonic resection or colostomy placement. The repair was accomplished with three steps: mucosal sealing of the perforation; an air leak test with the aid of local water irrigation (i.e. a water irrigation test); and direct observation for air bubbles during colonic air insufflation. Total serosal closure was completed with laparoscopic sutures.

Our approach with endoclips and laparoscopy provided a prompt repair of the colonic perforation and improved the outcome for the patient, who had an earlier discharge on the fourth postoperative day. ► **Table 1** lists four other cases [1–4] where endoclips have been used to treat colonic perforation.

Larger-scale studies are needed in order to determine whether using both laparoscopy and endoclips are superior to using endoclips alone.

Competing interests: None

Endoscopy_UCTN_Code_TTT_1AQ_2AG

Table 1 Salient clinical features of five cases described in the literature where endoclips were used to repair a colonic perforation.

Reference	Site of perforation	Time to recognition	Time to endoscopic surgery	Procedure	Discharge
Yoshikane et al. 1997	Descending colon	Immediate	Promptly	Endoclippping	14 days
Mana et al. 2001	Sigmoid colon	Immediate	Promptly	Endoclippping	> 1 week
Dhalla 2004	Cecum	Immediate	Promptly	Endoclippping	Not described
Barbagallo et al. 2007	Right flexure	Immediate	Promptly	Endoclippping	8 days
Senadhi et al. (December 2007; current report)	Sigmoid colon	Immediate	< 12 h for endoscopic and laparoscopic repair	Endoclippping and laparoscopic repair	4 days

V. Senadhi¹, A. Sharma¹, S. Fleischer²,
R. Franklin², S. Dutta³

¹ Johns Hopkins University/Sinai Hospital
Program in Internal Medicine, Sinai
Hospital, Baltimore, Maryland, USA

² Sinai Hospital of Baltimore, Baltimore,
Maryland, USA

³ Sinai Hospital of Baltimore and University
of Maryland School of Medicine,
Baltimore, Maryland, USA

References

- 1 Yoshikane H, Hidano H, Sakakibara A. Endoscopic repair by clipping of iatrogenic colonic perforation. *Gastrointest Endosc* 1997; 46: 464–466
- 2 Mana F, De Vogelaere K, Urban D. Iatrogenic perforation of the colon during diagnostic colonoscopy: endoscopic treatment with clips. *Gastrointest Endosc* 2001; 54: 258–259
- 3 Dhalla SS. Endoscopic repair of a colonic perforation following polypectomy using an endoclip. *Can J Gastroenterol* 2004; 18: 105–106
- 4 Barbagallo F, Castello G, Latteri S et al. Successful endoscopic repair of an unusual colonic perforation following polypectomy using an endoclip device. *World J Gastroenterol* 2007; 13: 2889–2891

Bibliography

DOI 10.1055/s-0030-1255718

Endoscopy 2010; 42: E213–E214

© Georg Thieme Verlag KG Stuttgart · New York ·
ISSN 0013-726X

Corresponding author

V. Senadhi

Johns Hopkins University/
Sinai Hospital Program in Internal Medicine
2606B Gage Court
Baltimore, MD 21209
USA

Fax: +1-678-623-5999

vsenadhi@hotmail.com