Primary lymphoma of the common bile duct presenting with acute pancreatitis and cholangitis

Malignant lymphoma primarily originating from the extrahepatic bile duct is extremely rare, and preoperative diagnosis is difficult because symptoms and imaging are nonspecific [1,2]. The most common symptoms of primary biliary lymphoma

are abdominal pain, weight loss, fever, and obstructive jaundice [1,2]. A total of 23 cases have been reported in literature [1–5], and, to the best of our knowledge, this is the fourth case of diagnosis made preoperatively and the first

report of primary biliary lymphoma with acute pancreatitis and cholangitis as the first clinical manifestations.

A 30-year-old male patient was referred to our hospital for acute pancreatitis. On admission he had severe abdominal pain, fever, and jaundice. Laboratory findings were: total leukocyte count, 15 700/ μ L; C-reactive protein, 7.89 mg/dL; aspartate aminotransferase, 271 U/L; alanine aminotransferase, 428 U/L; alkaline phosphatase, 580 U/L; γ -glutamyl transpeptidase, 428 U/L; serum total bilirubin, 9.1 mg/dL (direct 8.5 mg/dL); amylase, 951 U/L; and lipase, 1837 U/L. Transabdominal ultrasonography revealed only a dilated common bile duct (CBD) with sludge in the gallbladder. An urgent endoscopic ultrasound showed a stricture in the distal portion of the CBD, with proximal bile duct dilatation (Fig. 1).

During the same endoscopic session, endoscopic retrograde cholangiopancreatography (ERCP) and cholangiography confirmed the stricture in the lower portion of the CBD, with proximal dilatation (Fig. 2a); sphincterotomy and forceps biopsies were carried out (Fig. 2b), and finally a plastic stent was placed for drainage (Fig. 2c).

Histological examination (Fig. 3a) and immunohistochemistry (Fig. 3b–d) re-

Fig. 1 Endoscopic ultrasound (EUS) image of the stricture in the distal portion of the common bile duct, with proximal dilatation of the bile duct.

Fig. 2 Endoscopic retrograde cholangiopancreatography (ERCP) images of the stricture in the lower portion of the common bile duct: **a** proximal bile duct dilatation; **b** forceps biopsies; and **c** biliary stenting.

Fig. 3 **a** Histologic section and **b–d** immunohistochemistry of the endoscopic biopsy samples.

Fig. 4 Abdominal computed tomography (CT) images of the common bile duct mass with no metastasis.

vealed a large B-cell-type malignant lymphoma.

Total body computed tomography (CT) showed a well-circumscribed, heterogeneous, enhancing mass in the lower-mid portion of the CBD, without evidence of metastases (► **Fig. 4**).

The patient was symptom-free and scheduled for surgical resection. Examination of the surgical specimen confirmed the diagnosis of large B-cell-type malignant lymphoma, without metastasis in the lymph nodes included in the resection. The patient was referred to an oncologist for chemotherapy (rituximab, cyclophosphamide, vincristine, doxorubicin, and prednisolone [R-CHOP]) and after 6 months of follow-up he remains asymptomatic.

Competing interests: None

Endoscopy_UCTN_Code_CCL_1AZ_2AC

C. Luigiano¹, F. Ferrara¹, C. Fabbri¹, S. Ghersi¹, M. Bassi¹, A. M. Polifemo¹, P. Billi¹, A. Fornelli², F. Cinquantini³, N. D'Imperio¹

¹ Unit of Gastroenterology and Digestive Endoscopy, AUSL Bologna Bellaria-Maggiore Hospital, Bologna, Italy

² Unit of Anatomic Pathology, AUSL Bologna Maggiore Hospital, Bologna, Italy

³ Unit of Radiology, AUSL Bologna Maggiore Hospital, Bologna, Italy

References

- 1 Das K, Fisher A, Wilson DJ *et al*. Primary non-Hodgkin's lymphoma of the bile ducts mimicking cholangiocarcinoma. *Surgery* 2003; 134: 496–500
- 2 Joo YE, Park CH, Lee WS *et al*. Primary non-Hodgkin's lymphoma of the common bile duct presenting as obstructive jaundice. *J Gastroenterol* 2004; 39: 692–696
- 3 Sugawara G, Nagino M, Oda K *et al*. Follicular lymphoma of the extrahepatic bile duct mimicking cholangiocarcinoma. *J Hepatobiliary Pancreat Surg* 2008; 15: 196–199

4 Dote H, Ohta K, Nishimura R *et al*. Primary extranodal non-Hodgkin's lymphoma of the common bile duct manifesting as obstructive jaundice: report of a case. *Surg Today* 2009; 39: 448–451

5 Yoon MA, Lee JM, Kim SH *et al*. Primary biliary lymphoma mimicking cholangiocarcinoma: a characteristic feature of discrepant CT and direct cholangiography findings. *J Korean Med Sci* 2009; 24: 956–959

Bibliography

DOI 10.1055/s-0030-1255766

Endoscopy 2010; 42: E265–E266

© Georg Thieme Verlag KG Stuttgart · New York · ISSN 0013-726X

Corresponding author

C. Luigiano, MD

Unit of Gastroenterology and Digestive Endoscopy
AUSL Bologna Bellaria-Maggiore Hospital
Largo Nigrisoli 2
Bologna
Italy

Fax: +39-0516478967

carmeluigiano@libero.it