Endoscopic ultrasound-guided fine-needle aspiration for the diagnosis of retroperitoneal schwannoma

Schwannoma is a rare peripheral nerve sheath tumor that is difficult to diagnose by imaging features alone. Endoscopic ultrasound (EUS)-guided fine-needle aspiration (FNA) with specific immunohistochemical (IHC) staining may be the only tool to obtain a diagnostic sample from such lesions. There are only a few case reports describing EUS-FNA diagnosis of retroperitoneal Schwannoma [1,2]. In this report, we describe four cases (three males; mean age: 54.5 ± 16.4 years) with retroperitoneal Schwannoma, in whom the diagnosis was achieved with EUS-FNA and adjunctive IHC staining.

All lesions were well demarcated, and had a rounded contour (● Fig. 1 a).

The mean size of the lesions on EUS was 23.7 ± 3.6 mm. EUS-FNA was successfully performed with a 22-gauge needle in all cases (● Fig. 1 b), with a sufficient yield for both cytological and cellblock analysis. The median number of needle passes was 2.5 (range 2–3). The cellblock analysis revealed bland proliferation of spindle cells with a palisading appearance and wavy fibrillar architecture (● Fig. 1 c). Further evaluation with IHC revealed negative staining for actin, CD34, CD-117, and strong positive staining for S-100 antibody in all cases (● Fig. 1 d). Further evaluation of the cellular proliferative activity was studied with Ki-67 staining, and a low proliferation rate (Ki-67 < 5%) was reported in all cases, supporting the benign nature of the lesions.

We recommended conservative follow-up for our patients rather than surgical resection, because all of the patients were asymptomatic and there were no mitotic figures on FNA, with a low Ki-67 index in all the aspirates. It is worth noting that most reports have stressed on complete surgical resection as the appropriate management of retroperitoneal schwannomas [3,4]. Our view is that the morbidity associated with surgical resection is not justified in these benign lesions, and the use of EUS-FNA to establish the diagnosis may help in avoiding unnecessary surgery.

Competing interests: None

Endoscopy_UCTN_Code_CCL_1AF_2AG

Fig. 1 Imaging findings and histopathological features. **a** Contrast enhanced computed tomography (CT) scan showing well-demarcated low density mass in the retroperitoneal region. **b** Endoscopic ultrasound image showing a fine needle inserted into the mass (arrow). **c** Spindle cells on cellblock sections, without any mitosis (hematoxylin and eosin stain, original magnification × 400). **d** Immunohistochemical S-100 positive staining (magnification × 400).

S. Hijioka¹, A. Sawaki¹, N. Mizuno¹, K. Hara¹, M. A. Mekky², V. Bhatia³, W. Hosoda⁴, Y. Yatabe⁴, Y. Shimizu⁵, K. Tamada⁶, Y. Niwa⁷, K. Yamao¹

¹ Department of Gastroenterology, Aichi Cancer Center Hospital, Nagoya, Japan

² Department of Tropical Medicine and Gastroenterology, Assiut University Hospital, Assiut, Egypt

³ Department of Medical Hepatology, Institute of Liver and Biliary Sciences, Delhi, India

⁴ Department of Genetic Pathology and Molecular Diagnostics, Aichi Cancer Center Hospital, Nagoya, Japan

⁵ Department of Gastroenterological Surgery, Aichi Cancer Center Hospital, Nagoya, Japan

⁶ Department of Gastroenterology, Jichi Medical University, Tochigi, Japan

⁷ Department of Endoscopy, Aichi Cancer Center Hospital, Nagoya, Japan

References

- Okada N, Hirooka Y, Itoh A et al. Retroperitoneal neurilemoma diagnosed by EUS-guided FNA. *Gastrointest Endosc* 2003; 57: 790–792
- Facciorusso D, Federici T, Giacobbe A et al. Retroperitoneal neurilemoma diagnosed by endosonographically guided fine needle aspiration. *J Clin Ultrasound* 2006; 34: 241–243
- Li Q, Gao C, Juzi JT, Hao X. Analysis of 82 cases of retroperitoneal schwannoma. *ANZ J Surg* 2007; 77: 237–240
- Goh BK, Tan YM, Chung YF et al. Retroperitoneal schwannoma. *Am J Surg* 2006; 192: 14–18

Bibliography

DOI 10.1055/s-0030-1255786

Endoscopy 2010; 42: E296

© Georg Thieme Verlag KG Stuttgart · New York · ISSN 0013-726X

Corresponding author

S. Hijioka

Department of Gastroenterology

Aichi Cancer Center Hospital

1-1 Kanokoden

Chikusa-ku

Nagoya

Aichi 464-8681

Japan

Fax: +81-52-7642963

rizasusu@aichi-cc.jp