ERCP for the treatment of traumatic biliobronchial and biliocutaneous fistulas

Fig. 1 Fistulogram demonstrating pooling of contrast (white arrow) around the right costophrenic angle, communicating with the liver laceration. Multiple intrahepatic ducts are also visualized (black arrow).

A 21-year-old man presented to casualty following a roadside bomb blast. A traumatic pneumothorax necessitated the insertion of a chest drain, and a laparotomy revealed multiple liver lacerations and ureteric contusion. While the patient was awaiting further ureteric surgery after a percutaneous nephrostomy, the chest drain was removed. One week later, lime-green serosanguinous fluid began to leak from the drain site, while the patient began to cough up bitter yellow sputum. Biochemical analysis of these fluids con-

firmed the presence of bilirubin, suggesting the possibility of thoracobiliary fistulas involving the airways and pleural cavity. Ultrasound and computed tomography (CT) scans delineated a single biliocutaneous fistula, which was surgically excised. However, because the biliptysis and bile leakage persisted, a fistulogram (● Fig. 1) and contrast CT scan (● Fig. 2) were undertaken; these revealed multiple biliocutaneous and biliobronchial fistulas. Due to the patient's ongoing recovery from complex major surgery, a conservative approach involving endoscopic retrograde cholangiopancreatography (ERCP) with sphincterotomy and stenting was adopted. The ERCP findings were normal (i.e. revealing no contrast leakage), but the procedure resulted in the complete cessation of biliptysis and of cutaneous bile leakage (i.e. from 500 ml/day to nil) over 1 week. The patient thereafter remained fully asymptomatic from the thoracobiliary trauma.

Biliopleural and biliobronchial fistulas are hitherto unreported complications of blast injuries [1]. Regardless of etiology, the rare biliopleural and biliobronchial fistulas are diagnosed on finding bile in the pleural cavity, by thoracocentesis, or in a sputum sample [2,3]. Nonoperative fistula resolution may be attempted by distally decompressing the biliary system either surgically or endoscopically; the latter offers both diagnostic and therapeutic potential with a single minimally

traumatic procedure [4,5]. We recommend that even without evidence of biliary hypertension (i.e. if imaging excludes bile duct dilatation), ERCP with sphincterotomy and stenting is considered in patients like ours who have complex refractory pathology or are unfit for surgery. The relatively minor procedure may significantly reduce retrograde bile flow, resulting in major symptomatic improvement, before riskier alternatives (e.g. laparotomy or thoracotomy) are considered.

Endoscopy_UCTN_Code_TTT_1AR_2AG

Competing interests: None

S. G. Thrumurthy^{1,2}, A. H. M. P. Anurudha¹, M. I. M. De Zoysa¹, D. N. Samarakera¹

¹ University Surgical Unit, National Hospital of Sri Lanka, Colombo 10, Sri Lanka

² Department of General Surgery, Royal Preston Hospital, Lancashire Teaching Hospitals NHS Foundation Trust, Preston, PR2 9HT, UK

References

- 1 Wolf SJ, Beberta VS, Bonnett CJ et al. Blast injuries. *Lancet* 2009; 374: 405–415
- 2 Lewis JR, Te HS, Gehlbach B et al. A case of biliopleural fistula in a patient with hepatocellular carcinoma. *Nat Rev Gastroenterol Hepatol* 2009; 6: 248–251
- 3 Pende V, Marchese M, Mutignani M et al. Endoscopic management of biliopleural fistula and biloma after percutaneous radiofrequency ablation of liver metastasis. *Gastrointest Endosc* 2007; 66: 616–618
- 4 Chua HK et al. Bronchobiliary fistula: principles of management. *Ann Thorac Surg* 2000; 70: 1392–1394
- 5 Singh B, Moodley J, Sheik-Gafoor MH et al. Conservative management of thoracobiliary fistula. *Ann Thorac Surg* 2002; 73: 1088–1091

Bibliography

DOI 10.1055/s-0030-1255898

Endoscopy 2011; 43: E42

© Georg Thieme Verlag KG Stuttgart · New York · ISSN 0013-726X

Corresponding author

S. G. Thrumurthy, MBChB (Hons)

Department of General Surgery
Royal Preston Hospital
Lancashire Teaching Hospitals
NHS Foundation Trust
Preston
PR2 9HT
UK

Fax: +44-1257-245495
srigan@doctors.org.uk

Fig. 2 Contrast computed tomography (CT) scan of the abdomen demonstrating a biloma in the right hepatic lobe (white arrows) and one of many biliocutaneous tracts leading to a defect in the skin (black arrow). Air is demonstrated within the biloma (white arrowhead) as part of the tract.