Endoscopic treatment of acute ascending cholangitis in a patient with Roux-en-Y limb obstruction after a Whipple operation

Fig. 1 Computed tomography (CT) scan showed marked dilatation of the Roux-en-Y limb with a soft tissue mass, which was a suspected recurrent pancreatic carcinoma (white arrow).

Fig. 2 The endoscopic view revealed the obstructive point, past which the guidewire could be passed into the proximal part.

Fig. 3 Self-expandable metallic stent deployed under **a** fluoroscopic and **b** endoscopic controls.

A 60-year-old man was diagnosed with pancreatic head adenocarcinoma. He underwent pancreaticoduodenectomy (Whipple procedure). Four months later he developed obstructive jaundice, high fever, and chills. His total bilirubin level was 2.0 mg/dL (range 0.3–1.2 mg/dL) and alkaline phosphatase was 270 U/L (range 39–117 U/L). Computed tomography (CT) of the abdomen showed marked dilatation of the afferent limb and a suspected recurrent tumor in the pancreatic area (● Fig. 1).

The patient underwent an emergency endoscopy and a guidewire could be passed beyond the obstruction into the proximal part of the afferent limb (● Fig. 2). Endoscopic placement of a self-expandable metallic stent (enteral stent, 80 mm; Boston Scientific, Massachusetts, USA) was performed using a therapeutic gastroscope (1TGIF; Olympus Corp., Tokyo, Japan) (● Fig. 3). After the procedure, the patient's bilirubin normalized and the fever resolved. Plain film of the abdomen revealed a good deployment of the stent (● Fig. 4).

The patient was discharged 1 week later. He received adjunctive radiotherapy and at the 6-month follow-up remained asymptomatic.

Afferent loop obstruction is a rare complication after the Whipple procedure. In the rare case of complete obstruction, there is a high risk of developing necrosis and perforation. This condition requires immediate intervention. Percutaneous transhepatic drainage [1,2] and surgical drainage [3,4] are alternative management strategies with very high risk in some patients. Endoscopic metallic stent placement [5] is a treatment of choice to avoid an unnecessary and high risk operation. Long-term follow-up data are required to establish its clinical efficacy.

Competing interests: None

Endoscopy_UCTN_Code_TTT_1AR_2AK

Fig. 4 One day later, plain film of the abdomen demonstrated the metallic stent was placed in a good position.

T. Akaraviputh, A. Trakarnsanga, K. Tolan

Minimally Invasive Surgery Center,
Division of General Surgery, Faculty of
Medicine Siriraj Hospital, Mahidol
University, Bangkok, Thailand

References

- 1 Yoshida H, Mamada Y, Taniai N et al. One-step palliative treatment method for obstructive jaundice caused by unresectable malignancies by percutaneous transhepatic insertion of an expandable metallic stent. *World J Gastroenterol* 2006; 12: 2423–2426
- 2 Johnsson E, Delle M, Lundell L, Liedman B. Transhepatic placement of an enteral stent to treat jaundice in a tumor recurrence obstructed afferent loop after a Whipple procedure. *Dig Surg* 2003; 20: 329–331
- 3 Aimoto T, Uchida E, Nakamura Y et al. Malignant afferent loop obstruction following pancreaticoduodenectomy: report of two cases. *J Nippon Med Sch* 2006; 73: 226–230
- 4 Spiliotis J, Karnabatidis D, Vaxevanidou A et al. Acute cholangitis due to afferent loop syndrome after a Whipple procedure: a case report. *Cases J* 2009; 2: 6339
- 5 Burdick JS, Garza AA, Magee DJ et al. Endoscopic management of afferent loop syndrome of malignant etiology. *Gastrointest Endosc* 2002; 55: 602–605

Bibliography

DOI 10.1055/s-0030-1255978

Endoscopy 2010; 42: E335–E336

© Georg Thieme Verlag KG Stuttgart · New York ·
ISSN 0013-726X

Corresponding author

T. Akaraviputh, MD

Minimally Invasive Surgery Center
Division of General Surgery
Department of Surgery
Faculty of Medicine Siriraj Hospital
Mahidol University
Bangkok 10700
Thailand
Fax: +66-2-412-1370
sitak@mahidol.ac.th