Gastric rupture before puncture of the stomach in percutaneous gastrostomy

Fig. 1 No gross abnormalities were found when a flexible endoscope was inserted into the stomach.

Fig. 2 During gastric insufflations, performed before puncture of the stomach for a percutaneous endoscopic gastrostomy (PEG), several fusiform-shaped tears developed along the lesser curvature of the proximal stomach.

A 73-year-old man had swallowing difficulties because of a recent cerebrovascular accident, and was on enteral feeding via a nasogastric tube. Percutaneous

endoscopic gastrostomy (PEG) was chosen for long-term enteral feeding. Before the PEG tube was inserted, a flexible endoscope was inserted into the stomach for inspection. No gross abnormal lesion was detected in the upper gastrointestinal tract (● Fig. 1).

Then, before puncturing, the stomach was insufflated and indentation of the gastric lumen was confirmed by finger palpation of the abdominal wall. This caused the patient to choke a few times. Subsequently, bleeding occurred in the upper stomach. Several fusiform-shaped tears had developed along the lesser curvature of the proximal stomach (● Fig. 2).

Computed tomography (CT) imaging revealed pneumoperitoneum (● Fig. 3a) and pneumomediastinum (● Fig. 3b), requiring emergency laparotomy.

A 2-cm-long full-thickness tear along the lesser curvature of the stomach close to the cardia was identified and sutured.

Gastric rupture is caused by increased gastric pressure resulting from increased intra-abdominal pressure, with or without overdistention caused by food or gas. In more than 70% of the adult cases, gastric rupture occurs in the less distensible proximal lesser curvature of the stomach [1]. Spontaneous gastric rupture occurs due to vomiting, vigorous coughing, or convulsion. Its cause may also be iatrogenic, resulting from cardiopulmonary resuscitation, inadvertent esophageal intubation, Heimlich maneuver, or esophago-gastroduodenoscopy (EGD) [1–3].

Diagnostic EGD is extremely safe, and perforation of the gastrointestinal tract is rare with an incidence of 0.001%–0.05% [4]. Gastric rupture during gastric insuff-

lations, which are performed before puncturing the stomach for a PEG, is also rarely reported [5]. Endoscopic gastric insufflation with air is important to avoid colon injury. In the present case, the patient's choking during gastric insufflations, which led to a sudden increase in intra-abdominal pressure, may have caused the gastric rupture. Endoscopists should consider this rare yet potentially lethal complication.

Endoscopy_UCTN_Code_CPL_1AH_2AI

Competing interests: None

Y. Matsunaga¹, A. Goto¹, H. Kaneko², N. Hasegawa², H. Wakasugi¹, M. Itoh¹, K. Fujii¹, K. Suzuki¹, K. Nakahata¹, Y. Shinomura³

¹ Department of Gastroenterology, Kushiro City General Hospital, Kushiro, Japan

² Department of Surgery, Kushiro City General Hospital, Kushiro, Japan

³ First Department of Internal Medicine, Sapporo Medical University, Sapporo, Japan

References

- 1 Barker SJ, Karagianes T. Gastric barotrauma: a case report and theoretical considerations. *Anesth Analg* 1985; 64: 1026–1028
- 2 Tung PH, Law S, Chu KM et al. Gastric rupture after Heimlich maneuver and cardiopulmonary resuscitation. *Hepatogastroenterology* 2001; 48: 109–111
- 3 Ho CM, Yin IW, Tsou KF et al. Gastric rupture after awake fiberoptic intubation in a patient with laryngeal carcinoma. *Br J Anaesth* 2005; 94: 856–858
- 4 Misra T, Lalor E, Fedorak RN. Endoscopic perforation rates at a Canadian university teaching hospital. *Can J Gastroenterol* 2004; 18: 221–226
- 5 Schrag SP, Sharma R, Jaik NP et al. Complications related to percutaneous endoscopic gastrostomy (PEG) tubes. A comprehensive clinical review. *J Gastrointest Liver Dis* 2007; 16: 407–418

Bibliography

DOI 10.1055/s-0030-1256077

Endoscopy 2011; 43: E57

© Georg Thieme Verlag KG Stuttgart · New York · ISSN 0013-726X

Corresponding author

A. Goto, MD, PhD

Department of Gastroenterology
Kushiro City General Hospital
1-12, Shunkodai
Kushiro, 085-0822
Japan
Fax: +81-0154-414080
goa@sapmed.ac.jp

Fig. 3 Computed tomography (CT) imaging revealed a massive pneumoperitoneum and b pneumomediastinum.