EUS-guided choledochoduodenostomy for biliary drainage using tapered-tip plastic stent with multiple fangs

A 52-year-old man presented with abdominal pain and jaundice for 2 months. Computed tomography (CT) revealed a huge mass at the pancreatic head causing distal common bile duct (CBD) obstruction with superior mesenteric vein and superior mesenteric artery encasement (► Fig. 1).

The man underwent endoscopic retrograde cholangiopancreatography (ERCP), but we could not pass the duodenoscope through the duodenum because of tumor invasion. Therefore, a self-expandable metallic stent (SEMS) (Wallstent TM; Boston Scientific, Maryland, USA) was inserted. He underwent ERCP 2 weeks later but the ampulla was obscured. Therefore, endoscopic ultrasound (EUS) was considered for internal biliary drainage. The EUS showed a complex mass, 5.2×3.3 cm, at the pancreatic head, and the CBD was 2.05 cm (► Fig. 2).

After EUS-guided cholangiography, tailor-made Teflon dilators – 7 and 8.5 Fr – were used for dilation over the wire (► Figs. 3 and 4).

Then an 8.5 Fr $\times$ 6.5 cm tailor-made tapered-tip plastic stent, with multiple fangs but without a side hole, was inserted, and gave satisfactory drainage (► Figs. 5 and 6).

The patient was discharged without any complications. He was scheduled for SEMS insertion 4 months later.

In advanced pancreatic cancer, such as this case, percutaneous transhepatic biliary drainage (PTBD) and surgical drainage are the alternative options. PTBD is reported to have a higher complication rate of 10%–30%, while surgery is associated with a 2%–5% mortality and 17%–37% morbidity [1]. Even though EUS-guided biliary drainage was reported to be the safe and feasible procedure [2,3], it is not widely used because it requires more-advanced endoscopic skills. Possible complications of the EUS-guided biliary drainage, for example bile leakage and pneumoperitoneum, have also been reported. We minimized the leakage in this case by using a dilator instead of a needle knife or balloon dilation. We made the plastic stent ourselves instead of using a commercial one to make stent insertion easi-

Fig. 1 Computed tomography (CT) scan showing pancreatic mass at head and common bile duct, and pancreatic duct dilatation.

Fig. 2 Endoscopic ultrasound showed the common bile duct, 2.05 cm.

Fig. 3 Dilatation of the common bile duct.

Fig. 4 Cholangiogram.

Fig. 5 Our tailormade tapered-tip plastic stent with multiple fangs and no side hole.

Fig. 6 After stent insertion.

er, prevent bile leakage, and prevent CBD injury during stent insertion. Our idea of multiple fangs without a side-hole was to prevent stent migration and early clogging.

Endoscopy_UCTN_Code_TTT_1AS_2AD

Competing interests: None

V. Prachayakul^{1,2}, P. Aswakul², U. Kachintorn^{1,2}

¹ Division of Gastroenterology, Department of Internal Medicine, Siriraj Hospital, Mahidol University, Bangkok, Thailand

² Vikit Viranuwatti Endoscopy Center, Siriraj Hospital, Mahidol University, Bangkok, Thailand

References

- 1 van Delden OM, Lameris JS. Percutaneous drainage and stenting for palliation of malignant bile duct obstruction. *Eur Radiol* 2008; 18: 448–456
- 2 Savides TJ, Varadarajulu S, Palazzo L. EUS 2008 Working Group document: evaluation of EUS-guided hepaticogastrostomy. *Gastrointest Endosc* 2009; 69: S3–S6
- 3 Itoi T, Sofuni A, Itokawa F et al. Endoscopic ultrasonography-guided biliary drainage. *J Hepatobiliary Pancreat Surg* 2010; 17: 611–616

Bibliography

DOI 10.1055/s-0030-1256140

Endoscopy 2011; 43: E109–E110

© Georg Thieme Verlag KG Stuttgart · New York · ISSN 0013-726X

Corresponding author

P. Aswakul

Department of Internal Medicine

Siriraj Hospital

BangkokNoi

Bangkok 10711

Thailand

Fax: +66-2-4299672

asawakul@gmail.com