Successful management of aorto-esophageal fistula by combining endoscopic cyanoacrylate injection and endovascular stent grafting

Fig. 1 **a** Esophagogastroduodenoscopy (EGD) showing one 3-cm mass lesion with spurting bleeding 26 cm from incisors (arrow). **b** EGD after endovascular stent grafting, showing a submucosal mass measuring 3 cm × 1.8 cm with an erosive tip and *N*-butyl-2-cyanoacrylate coating (arrow).

Aorto-esophageal fistula (AEF) is a rare cause of catastrophic gastrointestinal bleeding. Thoracic aortic aneurysms are the leading cause of AEF [1]. Early diagnosis and prompt endovascular and/or surgical intervention before massive exsanguinating hemorrhage are key to survival [2]. Temporary endoscopic management with esophageal stent placement has been reported as a bridge therapy [3,4]. We describe a case of AEF with active bleeding noted during an endoscopic examination that was treated with injection of *N*-butyl-2-cyanoacrylate (NBCA, Histoacryl), followed by successful placement of an endovascular aortic stent graft.

A 69-year-old man with a history of ruptured diverticula that had been managed surgically presented to the emergency department with a 1-day history of chest pain and tarry stool, with hema-

Fig. 2 **a** Chest CT showing a 5-mm sacular aneurysm on the anterior wall of the mid descending thoracic aorta, consistent with aortic-esophageal fistula. A high-attenuation lesion in the mid esophagus abutting the aortic aneurysm is consistent with a deposition of *N*-butyl-2-cyanoacrylate with lipiodol (arrow). **b** Three-dimensional volume-rendered CT angiogram of thoracic aorta revealing a sacular aneurysm (arrow) protruding into the esophagus.

temesis and hematochezia on the morning of admission. His heart rate was 95 beats/min and his blood pressure was 214/111 mmHg. Laboratory data showed a rapid decline in hemoglobin (13.1 g/dL to 10.2 g/dL in 3 hours). Emergent esophagogastroduodenoscopy (EGD) revealed a 3-cm submucosal mass lesion with a spurting vessel 26 cm from the incisors (• Fig. 1 a).

Two milliliters of NBCA mixed with lipiodol was injected locally, achieving temporary hemostasis. Contrast-enhanced

computed tomography of the chest disclosed a descending thoracic aortic aneurysm with a 5-mm AEF (• Fig. 2).

Aorta angiography revealed a 5-mm aneurysm without active extravasation in the mid thoracic aorta (• Fig. 3 a).

The thoracic aortic aneurysm was repaired with a Cook Zenith II (34 mm × 77 mm) stent graft without complications (• Fig. 3 b). EGD after stent grafting the same day showed a submucosal mass measuring 3 cm × 1.8 cm with an erosive lesion and NBCA coating (• Fig. 1 b).

Fig. 3 **a** Aortogram showing a 5-mm aneurysm (arrow) from the mid thoracic aorta just below the carina without active extravasation into the esophagus. **b** After stent grafting, the aneurysm has disappeared.

After stent grafting, the patient suffered from intermittent episodes of chest and back pain. Intravenous broad-spectrum antibiotics were administered for a total of 8 weeks, followed by oral antibiotics. The patient recovered and remained healthy at 8 months follow-up.

Endoscopy_UCTN_Code_CCL_1AB_2AC_3AG

Competing interests: None

K.-C. Tseng^{1,2}, C.-W. Lin³, J. W.-H. Tan⁴

- ¹ Department of Internal Medicine, Buddhist Dalin Tzu Chi General Hospital, Chia-Yi, Taiwan
- ² School of Medicine, Tzuchi University, Hualien, Taiwan
- ³ Department of Radiology, Buddhist Dalin Tzu Chi General Hospital, Chia-Yi, Taiwan
- ⁴ Department of Cardiovascular Surgery, Buddhist Dalin Tzu Chi General Hospital, Chia-Yi, Taiwan

References

- 1 Léobon B, Roux D, Mugniot A *et al.* Endovascular treatment of thoracic aortic fistulas. *Ann Thorac Surg* 2002; 74: 247–249
- 2 Topel I, Stehr A, Steinbauer MG *et al.* Surgical strategy in aorto-esophageal fistulae: endovascular stentgrafts and in situ repair of the aorta with cryopreserved homografts. *Ann Surg* 2007; 246: 853–859
- 3 Park DH, Park JH, Lee SH *et al.* Temporary placement of a covered metal stent for the

management of a bleeding aorto-esophageal fistula. *Endoscopy* 2007; 39 (Suppl 1): E61–E62

- 4 Zuber-Jerger I, Hempel U, Rockmann F *et al.* Temporary stent placement in 2 cases of aorto-esophageal fistula. *Gastrointest Endosc* 2008; 68: 599–602

Bibliography

DOI 10.1055/s-0030-1256167

Endoscopy 2011; 43: E135–E136

© Georg Thieme Verlag KG Stuttgart · New York · ISSN 0013-726X

Corresponding author

J. W.-H. Tan, MD

Department of Cardiovascular Surgery
Buddhist Dalin Tzu Chi General Hospital
No. 2, Min-Sheng Road
Dalin Town
Chia-Yi
Taiwan, 622
Fax: +886-5-2648006
neugine@gmail.com