Serious complication following use of StomaphyX

Roux-en-Y gastric bypass (RYGB) is a popular surgical procedure to treat morbid obesity [1]. StomaphyX (endoGastric Solutions Inc., Redmond, Washington, USA) is designed for incisionless transoral endoscopic plication and revision of the gastric pouch (EPRGP) in case of pouch dilation with weight gain [2,3]. Nonresorbable polypropylene fasteners create full-thickness plications of the pouch wall, reducing its volume. StomaphyX was cleared by the Food and Drug Administration (FDA) in April 2007 [4] and clinical results are becoming available, with only 3% early complications [5].

We present the case of a 38-year-old woman who underwent StomaphyX EPRGP in 2007, after RYGB in 2000. The procedure was carried out in a peripheral hospital by a surgeon and was not planned within the regulations governing ongoing clinical trials. After 3 days, the patient was referred to our hospital with fever and thoracic pain. Thoracic X-ray showed left lung empyema (● Fig. 1). Antibiotics were started and she underwent two thoracic surgical procedures to drain the left hemithorax and to remove the pleural membranes (● Fig. 2). Upper gastrointestinal endoscopy showed three StomaphyX fasteners through the distal oesophageal wall and a dilated gastric pouch with another three fasteners around the gastrojejunostomy (● Fig. 3). Because of persistent empyema, a third draining thoracotomy was carried out 14 days later, which revealed adhesions at the distal esophagus. *Streptococcus viridans* and non-aureus *Staphylococcus* were cultured and antibiotics started. After a 28-day stay in the hospital, the patient was fit enough to be discharged. At present, 4 years after the complicated StomaphyX procedure, the patient is still under treatment for postoperative neuralgic pain of the left hemithorax. She was not covered by a clinical trial insurance. This case report illustrates the risk of serious complications of StomaphyX and the ethical implications of using new devices outside the control of regulated clinical trials.

Endoscopy_UCTN_Code_CPL_1AH_2AJ

Competing interest: None

Fig. 1 Chest X-ray showing left lung empyema with a thoracic drain.

Fig. 2 Thoracic computed tomography (CT) scan showing empyema and mediastinitis with a surgical thoracic drain.

Fig. 3 Endoscopic image of the gastric pouch showing inadequate placement of fasteners around the gastrojejunostomy.

J. I. Lenz¹, G. J. Hubens², P. A. Pelckmans¹, T. G. Moreels¹

¹ Division of Gastroenterology and Hepatology, Antwerp University Hospital, Antwerp, Belgium

² Division of Abdominal, Pediatric and Reconstructive Surgery, Antwerp University Hospital, Antwerp, Belgium

References

- 1 Thompson CC, Slattery J, Bundga ME, Lautz DB. Peroral endoscopic reduction of dilated gastrojejunal anastomosis after Roux-en-Y gastric bypass: a possible new option for patients with weight regain. *Surg Endosc* 2006; 20: 1744–1748
- 2 Elder KA, Wolfe BM. Bariatric surgery: a review of procedures and outcomes. *Gastroenterology* 2007; 132: 2253–2271
- 3 Mikami D, Needleman B, Narula V et al. Natural orifice surgery: initial US experience utilizing the StomaphyX device to reduce gastric pouches after Roux-en-Y gastric bypass. *Surg Endosc* 2010; 24: 223–228
- 4 Gibson S. Endoscopic solutions receives FDA clearance for first transoral surgical product: StomaphyXTM gets clearance first (April 2, 2007). Available from <http://www.egseurope.com> Accessed: November 15, 2010
- 5 Leitman IM, Virk CS, Avgerinos DV et al. Early results of trans-oral endoscopic plication and revision of the gastric pouch and stoma following Roux-en-Y gastric bypass surgery. *JSL* 2010; 14: 217–220

Bibliography

DOI 10.1055/s-0030-1256258

Endoscopy 2011; 43: E173–E174

© Georg Thieme Verlag KG Stuttgart · New York · ISSN 0013-726X

Corresponding author

T. Moreels

Antwerp University Hospital

Division of Gastroenterology and Hepatology

Wilrijkstraat 10

B-2650 Antwerp

Belgium

Fax: +32-3-8214478

tom.moreels@uza.be