Subcapsular hepatic hematoma following ERCP: case report and review

A 76-year-old woman, who was receiving anticoagulation for atrial fibrillation, was referred to our center for management of a common bile duct stone diagnosed by endoscopic ultrasonography. Anticoagulation was suspended and endoscopic retrograde cholangiopancreatography (ERCP) was subsequently performed. Cannulation of the main bile duct with a 0.035-inch guide wire was achieved without complications. Endoscopic biliary sphincterotomy was performed and stone extraction with a Fogarty catheter was achieved successfully, without apparent complications. Subsequently, the patient developed sharp right upper quadrant pain 6 hours after the procedure, but showed no signs of hemodynamic instability, and laboratory data did not show any evidence of complications. By 24-hours after the procedure, she was asymptomatic and was discharged after the reintroduction of anticoagulation.

The patient consulted again 5 days later because of persistent pain. Abdominal examination elicited mild right upper quadrant pain without tenderness. Laboratory data showed hemoglobin 9.6 g/dL (normal range 12–15 g/dL) and hematocrit 30.7% (normal range 36–41%). Computed tomography showed two high-density collections consistent with hematomas within the subdiaphragmatic and subhepatic spaces (● Fig. 1). The patient was managed conservatively. Anticoagulation was discontinued and a broad-spectrum antibiotic (piperacillin–tazobactam) was administered. The patient was discharged 15 days after the ERCP, without any further complications.

Subcapsular hepatic hematoma is a rare complication of ERCP. There are few published reports of this unusual complication [1–10], which may be explained by accidental puncture of the intrahepatic biliary tree by the guide wire. In this case, the patient probably developed an initial hematoma 6 hours after the procedure, which worsened because of the resumption of anticoagulation.

From the literature [1–10] (● Table 1), there is unanimous concern about the risk of infection in these patients, and in all cases, except two where no detail was given, patients were treated with anti-

Fig. 1 Computed tomography showing subdiaphragmatic and subhepatic hematomas.

Table 1 Patient characteristics from the reports of hematoma post endoscopic retrograde cholangiopancreatography (ERCP).

	Age/ Sex	Indication for ERCP	Guide wire	Puncture	Treatment	Antibiotic
Ortega et al. 2000 [1]	81/M	Common bile duct stone	NA	Yes, positive culture	Percutaneous drainage	Yes
Horn et al. 2004 [2]	88/F	Pancreatic cyst	Yes	No	Observation	Yes
Chi et al. 2004 [3]	43/F	Pancreatic cancer	Yes	No	Embolization	Yes
Priego et al. 2007 [4]	30/F	Obstructive jaundice	NA	No	Surgery, positive culture	Yes
Petit-Laurent et al. 2007 [5]	98/M	Common bile duct stone	Yes	Yes, negative culture	Percutaneous drainage	NA
Bhati et al. 2007 [6]	51/F	Common bile duct stone	Yes	Yes, negative culture	Percutaneous drainage	NA
McArthur et al. 2008 [7]	71/M	Common bile duct stone	Yes	No	Observation	Yes
De la Serna et al. 2008 [8]	71/F	Common bile duct stone	Yes	Yes, negative culture	Observation	Yes
Cárdenas et al. 2008 [9]	54/M	Bile leak post liver trans- plant	Yes	No	Observation	Yes
Revuelto et al. 2010 [10]	41/M	Common bile duct stone	NA	No	Observation	Yes
Current report	76/F	Common bile duct stone	Yes	No	Observation	Yes

M, male; F, female; NA, details not available.

biotics. Most of the patients including our own (6/11) were observed; three were treated by percutaneous drainage; and one each by embolization and surgery. There were no long term complications.

Endoscopy_UCTN_Code_TTT_1AU_2AC

Competing interests: None

**D. Del Pozo¹, I. Moral¹, E. Poves¹,
C. Sanz¹, M. Martín²**

¹ Gastroenterology Unit, Hospital Universitario Príncipe de Asturias, Madrid, Spain

² Internal Medicine Department, Hospital Universitario Príncipe de Asturias, Madrid, Spain

References

- 1 Ortega Deballon P, Fernández Lobato R, García Septiem J et al. Liver hematoma following endoscopic retrograde cholangiopancreatography (ERCP). *Surg Endosc* 2000; 14: 767
- 2 Horn TL, Peña LR. Subcapsular hepatic hematoma after ERCP: case report and review. *Gastrointest Endosc* 2004; 59: 594–596
- 3 Chi KD, Waxman I. Subcapsular hepatic hematoma after guidewire injury during endoscopic retrograde cholangiopancreatography: management and review. *Endoscopy* 2004; 36: 1019–1021
- 4 Priego P, Rodríguez G, Mena A et al. Subcapsular liver hematoma after ERCP. *Rev Esp Enferm Dig* 2007; 99: 53–54
- 5 Petit-Laurent F, Scalone O, Penigaud M, Barbeys J. Subcapsular hepatic hematoma after endoscopic retrograde cholangiopancreatography: case report and literature review. *Gastroenterol Clin Biol* 2007; 31: 750–752
- 6 Bhati CS, Inston N, Wigmore SJ. Subcapsular intrahepatic hematoma: an unusual complication of ERCP. *Endoscopy* 2007; 39 (Suppl 1): E150
- 7 McArthur KS, Mills PR. Subcapsular hepatic hematoma after ERCP. *Gastrointest Endosc* 2008; 67: 379–380
- 8 De La Serna-Higuera C, Fuentes Coronel A, Rodríguez Gómez SJ, Martín Arribas MI. Subcapsular hepatic hematoma secondary to the use of hydrophilic guidewires during endoscopic retrograde cholangiopancreatography. *Gastroenterol Hepatol* 2008; 31: 266–267
- 9 Cárdenas A, Crespo G, Balderramo D et al. Subcapsular liver hematoma after Endoscopic Retrograde Cholangiopancreatography in a liver transplant recipient. *Ann Hepatol* 2008; 7: 386–388
- 10 Revuelto Rey J, Gordillo Escobar E, Batalha P. Subcapsular hepatic hematoma after ERCP. *Med Intensiva* 2010; 34: 224

Bibliography

DOI 10.1055/s-0030-1256267

Endoscopy 2011; 43: E164–E165

© Georg Thieme Verlag KG Stuttgart · New York · ISSN 0013-726X

Corresponding author

D. Pozo Prieto, MD

Aparato Digestivo

Hospital Universitario Príncipe de Asturias,

Alcalá de Henares,

28805 Madrid,

Spain

Fax: +34-91-8801825

dpozo.hupa@salud.madrid.org