Spontaneous tension pneumocephalus after esophagogastroscope

An 80-year-old man presented with generalized weakness and upper abdominal discomfort. A computed tomography (CT) brain scan showed hyperpneumatized paranasal sinuses but was otherwise normal (● Fig. 1).

An esophagogastroscope was performed which revealed gastritis. He received intravenous sedation during the procedure but was noted to have been gagging. He lapsed into coma 5 days later due to a massive tension pneumocephalus, which was urgently relieved through a burr hole (● Fig. 2).

A cisternography demonstrated a cerebrospinal fluid (CSF) fistula at the left frontal sinus. Surgical repair was performed through a burr hole, which confirmed the presence of the fistula and a bony defect on the posterior sinus wall (● Fig. 3). The patient recovered well.

Neurological complications of gastrointestinal endoscopy are extremely rare; cerebral air embolism has been described, but the occurrence of pneumocephalus has never been reported [1]. Pneumocephalus is an uncommon but potentially fatal condition. The majority of cases are traumatic in origin [2]. Spontaneous pneumocephalus may result from actions which generate high pressure within the paranasal sinuses, such as Valsalva's manoeuvre [3]. The presence of hyperpneumatized paranasal sinuses may also predispose to spontaneous pneumocephalus [4]. In the present case, a sudden rise in airway pressure during endoscopy, albeit transient, was likely to have resulted in the formation of a CSF fistula through a hyperpneumatized sinus. The resultant dural tear acted as a ball valve which allowed continuous inflow of air, and presented with the delayed onset of spontaneous pneumocephalus.

Spontaneous pneumocephalus may resolve on conservative treatment. Surgical treatment is indicated when there is evidence of raised intracranial pressure, neurological deterioration, or when the dural defect does not heal satisfactorily. We have successfully repaired the CSF fistula through a craniotomy, although endoscopic frontal outflow tract obliteration may be considered as a viable, minimally invasive alternative [5].

Fig. 1 Plain computed tomography (CT) scan showing hyperpneumatized air sinuses.

Fig. 2 Pneumocephalus after esophagogastroscope.

Endoscopy_UCTN_Code_CPL_1AH_2AJ

Competing interests: None

G. K. K. Leung

Division of Neurosurgery, Department of Surgery, Li Ka Shing Faculty of Medicine, The University of Hong Kong, Queen Mary Hospital, Hong Kong

Fig. 3 Cerebrospinal fluid fistula on the posterior wall of the left frontal sinus.

References

- 1 McAree BJ, Gilliland R, Campbell DM et al. Cerebral air embolism complicating esophagogastroduodenoscopy (EGD). *Endoscopy* 2008; 40 (Suppl 2): E191 – E192
- 2 Markham JW. The clinical features of pneumocephalus based upon a survey of 284 cases with report of 11 additional cases. *Acta Neurochir (Wien)* 1967; 16: 1 – 78
- 3 Schrijver HM, Berendse HW. Pneumocephalus by Valsalva's maneuver. *Neurology* 2003; 60: 345 – 346
- 4 Lee JS, Park YS, Kwon JT, Suk JS. Spontaneous pneumocephalus associated with pneumosinus dilatans. *J Korean Neurosurg Soc* 2010; 47: 395 – 398
- 5 Sindwani R. Endoscopic frontal outflow tract obliteration for pneumocephalus after frontal sinus cranialization. *Otolaryngol Head Neck Surg* 2008; 139: 735 – 737

Bibliography

DOI 10.1055/s-0030-1256392

Endoscopy 2011; 43: E208

© Georg Thieme Verlag KG Stuttgart · New York · ISSN 0013-726X

Corresponding author

G. K. K. Leung, MD

Division of Neurosurgery
Department of Surgery
Li Ka Shing Faculty of Medicine
The University of Hong Kong
Queen Mary Hospital
102 Pokfulam Road
Hong Kong
Fax: +852-28184350
gilberto@hkucc.hku.hk