Cholangioscope-assisted percutaneous fibrin glue sealing of bile leak complicating transarterial chemoembolization of hepatocellular carcinoma after liver transplantation

Fig. 1 Computed tomography (CT) scan showing: **a** biloma in hepatic segment VII with all-purpose drainage loop (APDL) indwelling catheter; **b**, **c** the fistula is well visualized between the biloma and the segmental biliary branch (black arrows).

Transarterial chemoembolization (TACE) is recommended for patients with unresectable hepatocellular carcinoma (HCC); however, it is not a risk-free procedure and biloma may occur as a complication [1].

A 45-year-old man, following liver transplantation, presented with recurrent HCC

Fig. 2 Percutaneous cholangioscopy confirming the biliary fistula.

Fig. 3 Cholangioscopic view of the fistula orifice.

Fig. 4 Cholangioscopy-assisted fibrin glue injections using a 19-G needle.

in the caudate lobe, close to the caval vein, and was treated by TACE. Subsequently, the patient was admitted for abdominal pain and fever. Computed tomography (CT)-guided percutaneous cholangiography confirmed the diagnosis of infected biloma (• **Figs. 1, 2**). During the following weeks there was abundant drainage, despite both external drainage and endoscopic treatment. It was decided to attempt direct closure of the fistula with a cholangioscope-assisted procedure. Briefly, an inverse rendezvous procedure was successfully carried out, allowing the retrieval of the endoscopic guide wire, fol-

lowed by insertion of a percutaneous wire-guided cholangioscope (Polyscope, Lumenis Inc., Santa Clara, California, USA) into the biloma. An angiographic introducer was inserted beside the cholangioscope and a 19-G needle was inserted in the introducer. The cholangioscopic approach allowed multiple fibrin glue injections (Tissucol, Baxter Healthcare, Deerfield, Illinois, USA) around the distal opening of the peripheral bile duct, for a total volume of 3 mL (• **Fig. 3–5**). A CT scan taken after a few days showed absence of fluid in the biloma, confirming healing of the biliary fistula (• **Fig. 6**).

Fig. 5 Cholangioscopic view after fibrin glue sealing of bile leak.

Fig. 6 Follow-up computed tomography (CT) scan showing the absence of fluid in the biloma in spite of the closure of the all-purpose drainage loop (APDL) catheter, thus demonstrating healing of the biliary fistula.

Conservative management of biloma allows resolution in more than 80% of cases [2–4]. However, cases resistant to well-established conservative strategies still represent a challenge. To our knowledge, this is the first report of a novel technique in the management of hepatic biloma. Use of fibrin glue injection to seal a bile leak could represent an indication for therapeutic cholangioscopy, although it requires confirmation through application in further patients.

Video 1

Cholangioscope-assisted percutaneous fibrin glue sealing of bile leak complicating transarterial chemoembolization of recurrent hepatocellular carcinoma after liver transplantation.

Endoscopy_UCTN_Code_TTT_1AR_2AG

Competing interests: None

V. Cennamo¹, L. Fuccio¹, E. Giampalma², E. Terzi³, L. H. Eusebi¹, C. Mosconi², F. Piscaglia³

¹ Division of Gastroenterology, Department of Digestive Disease and Internal Medicine, S. Orsola-Malpighi General and University Hospital, Bologna, Italy

² Division of Radiology, Department of Digestive Disease and Internal Medicine, S. Orsola-Malpighi General and University Hospital, Bologna, Italy

³ Division of Internal Medicine, Department of Digestive Disease and Internal Medicine, S. Orsola-Malpighi General and University Hospital, Bologna, Italy

References

- 1 Sakamoto I, Aso N, Nagaoki K *et al.* Complications associated with transcatheter arterial embolization for hepatic tumors. *RadioGraphics* 1998; 18: 605–619
- 2 Morelli J, Mulcahy HE, Willner IR *et al.* Endoscopic treatment of postliver transplantation biliary leaks with stent placement across the leak site. *Gastrointest Endosc* 2001; 54: 471–475
- 3 Kim JH, Ko GY, Sung KB *et al.* Early post-transplant hepatic venous outflow obstruction: Long-term efficacy of primary stent placement. *Liver Transplant* 2008; 14: 1142–1149
- 4 Shah JN. Endoscopic treatment of bile leaks: current standards and recent innovations. *Gastrointest Endosc* 2007; 65: 1069–1072

Bibliography

DOI 10.1055/s-0030-1256512

Endoscopy 2011; 43: E238–E239

© Georg Thieme Verlag KG Stuttgart · New York · ISSN 0013-726X

Corresponding author

V. Cennamo

Division of Gastroenterology
Department of Digestive Diseases
and Internal Medicine
S. Orsola-Malpighi General and University Hospital
40138 Bologna
Italy
Fax: +39-51-6363338
cennamoit@yahoo.it