Peroral cholangioscopy in a patient with a Billroth II gastrectomy using the SpyGlass Direct Visualization System

Peroral cholangioscopy in patients with altered gastrointestinal anatomy is challenging because of the technical difficulties associated with accessing the papilla and the high rate of complications experienced when using a side-viewing duodenoscope [1]. The recently developed SpyGlass Direct Visualization System (Microvasive Endoscopy, Boston Scientific, Natick, Massachusetts, USA) can be passed through a forward-viewing endoscope because its diameter is 10 Fr [2–4]. We describe a case of cholangiocarcinoma that was successfully visualized using a SpyGlass cholangioscope in a patient with a Billroth II gastrectomy.

An 81-year-old man presented to our institution for evaluation of obstructive jaundice. His medical history included a Billroth II gastrectomy for gastric cancer 10 years previously. A front-viewing endoscope was passed through the afferent limb reaching the papilla without difficulty. A cholangiogram showed a severe stricture at the lower end of the bile duct (● Fig. 1).

Following endoscopic papillary balloon dilation, a SpyGlass cholangioscope was passed into the bile duct without difficulty (● Fig. 2).

Cholangioscopy showed an irregular, fine granular lesion at the upper end of the stricture and normal mucosa at the hilar bile duct (● Fig. 3), both of which were biopsied under cholangioscopic imaging using a 3-Fr SpyBite forceps (Boston Scientific; ● Video 1).

The biopsy specimens taken from the irregular lesion showed adenocarcinoma, whereas those from the hilar bile duct showed no malignancy. He was diagnosed as having a lower bile duct carcinoma without intraepithelial spread of tumor to the hilar bile duct and underwent pancreaticoduodenectomy at our hospital.

Peroral cholangioscopy with mapping biopsies has been shown to be effective for the diagnosis of intraepithelial tumor spread in cholangiocarcinoma [5]. Although SpyGlass cholangioscopy through a colonoscope has been reported in patients with a Roux-en-Y anastomosis [3,4], to our knowledge, this is the first case of bile duct biopsy under direct visualization

Fig. 1 Cholangiogram showing a stricture at the lower end of the bile duct.

Fig. 2 Fluoroscopy showing the SpyGlass cholangioscope inside the bile duct, having been passed through the forward-viewing endoscope.

Fig. 3 Cholangioscopy showing: **a** an irregular, fine granular lesion at the upper end of the stricture; **b** normal mucosa at the hilar bile duct.

Endoscopy_UCTN_Code_TTT_1AR_2AB

Competing interests: None

K. Kawakubo, H. Isayama, T. Tsujino, Y. Nakai, N. Sasahira, H. Kogure, T. Sasaki, N. Yamamoto, K. Hirano, M. Tada, K. Koike

Department of Gastroenterology, Graduate School of Medicine, The University of Tokyo, Japan

Video 1

SpyGlass cholangioscopy showing the lesion at the upper end of the stricture and the normal hilar bile duct followed by a biopsy being taken with the SpyBite forceps.

References

- 1 *Hintze RE, Adler A, Veltzke W, Abou-Rebyeh H.* Endoscopic access to the papilla of Vater for endoscopic retrograde cholangiopancreatography in patients with billroth II or Roux-en-Y gastrojejunostomy. *Endoscopy* 1997; 29: 69–73
- 2 *Chen YK, Pleskow DK.* SpyGlass single-operator peroral cholangiopancreatography system for the diagnosis and therapy of bile-duct disorders: a clinical feasibility study (with video). *Gastrointest Endosc* 2007; 65: 832–841
- 3 *Mou S, Waxman I, Chennat J.* Peroral cholangioscopy in Roux-en-Y hepaticojejunostomy anatomy by using the SpyGlass Direct Visualization System (with video). *Gastrointest Endosc* 2010; 72: 458–460
- 4 *Baron TH, Saleem A.* Intraductal electrohydraulic lithotripsy by using SpyGlass cholangioscopy through a colonoscope in a patient with Roux-en-Y hepaticojejunostomy. *Gastrointest Endosc* 2010; 71: 650–651
- 5 *Kawakami H, Kuwatani M, Etoh K et al.* Endoscopic retrograde cholangiography versus peroral cholangioscopy to evaluate intraepithelial tumor spread in biliary cancer. *Endoscopy* 2009; 41: 959–964

Bibliography

DOI 10.1055/s-0030-1256606

Endoscopy 2011; 43: E241–E242

© Georg Thieme Verlag KG Stuttgart · New York ·
ISSN 0013-726X

Corresponding author

K. Kawakubo, MD

Department of Gastroenterology,
Graduate School of Medicine

The University of Tokyo

7-3-1 Hongo Bunkyo-ku

Tokyo 113-8655

Japan

Fax: +81-33-8140021

kkawakubo-gi@umin.ac.jp