Arytenoid dislocation following upper gastrointestinal endoscopy

A 46-year-old female teacher presented for upper gastrointestinal endoscopy. A standard Olympus endoscope (GIF H190, Olympus, Center Valley, Pennsylvania, USA) was used for evaluation of gastritis. After 12 hours, the patient developed hoarseness, throat pain, and swelling. A computed tomography (CT) scan, barium swallow, and laryngeal electromyography (LEMG) studies showed normal findings. Stroboscopy showed a higher left vocal cord with axis deviation and persistent glottal gap of 1 mm (Fig. 1). After 2 months the patient underwent laryngoplasty with micronized dermis injection and reduction of the left arytenoid dislocation. Dysphonia gradually improved over 6 months with voice therapy. Signs and symptoms of arytenoid dislocation include hoarseness, breathiness, vocal fatigue, aphonia, as well as dysphagia. Diabetes mellitus and renal failure can weaken the arytenoid joint [1], and use of airway tools such as a misplaced laryngoscope, laryngeal mask airways [2] and transesophageal echocardiography (TEE) probe [3] has also resulted in arytenoid dislocations. In the present case, the initial insertion of the gastroscope was not carried out under direct visualization, thus resulting in traumatic arytenoid dislocation. An unrecognized cricoarytenoid joint dislocation is often mistaken for vocal fold paralysis, and treatment is delayed. Direct laryngoscopy and CT can be useful in the diagnosis of patients with arytenoid dislocation. LEMG evaluates innervation of the laryngeal muscles, distinguishing between paralysis and dislocation. An invaluable tool for diagnosis is stroboscopy [4], which provides a magnified slow-motion view of the vocal cords. Successful treatment is frequently predicated on early intervention. Voice therapy is only indicated when hoarseness has an etiologic diagnosis and is an important

Fig. 1 a Rigid endoscopy during adduction showing asymmetric arytenoid closure with posterior gap in a patient with hoarseness, throat pain, and swelling. b Rigid endoscopy during inspiration showing the lateralized arytenoid.

adjunct for patients [5]. Surgical correction is the treatment of choice, with botulinum toxin injections for laryngeal rebalancing. Gastroenterologists must add arytenoid dislocation to the list of potential complications that can occur with gastrointestinal endoscopy. Patients complaining of hoarseness or other neck symptoms after an upper gastrointestinal endoscopy should undergo upper airway evaluation for early diagnosis and treatment.

Endoscopy_UCTN_Code_CPL_1AH_2AJ

Competing interests: Dr. P. Woo is a compensated speaker for LifeCell Corporation.

A. Afonso¹, P. Woo², A. Reed¹

¹ Department of Anesthesiology, Mount Sinai Hospital, New York, USA

² Department of Otolaryngology, Mount Sinai Hospital, New York, USA

Reference

- 1 Quick CA, Merwin GE. Arytenoid dislocation. *Arch Otolaryngol* 1978; 104 (5): 267–270
- 2 Cros AM, Pitti R, Conil C et al. Severe dysphonia after use of a laryngeal mask airway. *Anesthesiology* 1997; 86 (2): 498–500
- 3 Niwa Y, Nakae A, Ogawa M et al. Arytenoid dislocation after cardiac surgery. *Acta Anaesthesiol Scand* 2007; 51 (10): 1397–1400
- 4 Sataloff RT, Spiegel JR, Hawkshaw MJ. Stroboscopy: Results and clinical value. *Ann Otol Rhinol Laryngol* 1991; 100 (9 Pt 1): 725–727
- 5 Mau T. Diagnostic evaluation and management of hoarseness. *Med Clin North Am* 2010; 94 (5): 945–960

Bibliography

DOI 10.1055/s-0030-1256690

Endoscopy 2011; 43: E368

© Georg Thieme Verlag KG Stuttgart · New York · ISSN 0013-726X

Corresponding author

A. Afonso

Mount Sinai Medical Center
Department of Anesthesiology
One Gustave L. Levy Place
Box 1010
New York
NY 10029-6574
USA
Fax: +212-426-2009
Anoushka.Afonso@mssm.edu