Hematemesis from ruptured aberrant right hepatic artery aneurysm eroding through the duodenal wall

Fig. 1 Large submucosal mass with ulcer in the duodenal bulb.

Fig. 2 An oral and intravenous contrast-enhanced computed tomography (CT) scan demonstrates a pseudoaneurysm (red arrow) with a surrounding hematoma (green arrows) just above the duodenal bulb and medial to the left lobe of the liver.

A 56-year-old man presented with hematemesis and multiple episodes of melena. He had a history of chronic lymphocytic leukemia and traumatic rupture of the spleen leading to splenectomy and splenic artery embolization 6 years earlier. Esophagogastroduodenoscopy (EGD) revealed a large submucosal mass (7 × 5 cm) with an ulcerated overlying area associated with clot in the duodenal bulb (• **Fig. 1**). The ulcer was treated with epinephrine (1:10000). A computed tomography (CT) scan of the abdomen revealed a 2.5 × 1.8-cm pseudoaneurysm from an aberrant hepatic artery off the superior mesenteric artery (• **Fig. 2**) along with surrounding hematoma, causing mass effect on the duodenum; this was further confirmed with a CT angiogram (• **Fig. 3a**).

Coil embolization was performed with complete obliteration of the hepatic artery pseudoaneurysm (• **Fig. 3b**). The patient was subsequently discharged home after 4 days of observation.

The patient presented 2 months later with recurrent episodes of melena. A CT angiogram showed no active extravasations. EGD revealed a long segment of coil protruding from the pylorus into the stomach, along with coffee ground materials. There was a large mound-like focal bulge at the superior aspect of the duodenal bulb, with a 6-mm defect without active bleeding, along with the protruding coil (• **Fig. 4**). The patient underwent a distal gastrectomy, Billroth II gastrojejunos-

Fig. 3 **a** An angiogram performed immediately after the computed tomography (CT) scan confirms the pseudoaneurysm (red arrow) arising from a replaced hepatic artery (white arrow) off the superior mesenteric artery. **b** Post-embolization arteriogram of the common hepatic artery, demonstrating occlusion of the two large pseudoaneurysms.

Fig. 4 Long segment of coil protruding from the pylorus into the stomach.

tomy, and ligation of gastroduodenal artery. He was discharged after 5 days of observation and remained well without further episodes of bleeding after 6 months of follow-up.

Hepatic artery pseudoaneurysm is a rare cause of upper gastrointestinal bleeding, and can be life-threatening [1,2]. Angiographic embolization is an effective method of treatment with a reported success rate of 80–100% [3]. However, complications from embolization are not unusual, as noted in our case with extrusion of coils through the duodenal wall with potential for re-bleeding. Surgery may be needed in unusual circumstances for more definitive therapy.

Endoscopy_UCTN_Code_CCL_1AB_2AZ_3AD

Competing interests: None

Y. J. Kim¹, S. K. Satapathy², L. Law²,
A. Volfson², B. Friedman³, S. Yang²,
C. Sung³, D. S. Siegel³, B. DeVito⁴

¹ Department of Internal Medicine,
Hofstra North Shore-LIJ Health system
at Long Island Jewish Medical Center,
New York, USA

² Division of Gastroenterology, Hofstra
North Shore-LIJ Health system at Long
Island Jewish Medical Center, New York,
USA

³ Department of Radiology, Hofstra North
Shore-LIJ Health system at Long Island
Jewish Medical Center, New York, USA

⁴ Division of Gastroenterology, Hofstra
North Shore-LIJ Health system at North
Shore University Hospital, New York, USA

References

- 1 Poon R, Tuen H, Yeung C *et al.* GI haemorrhage from fistula between right hepatic artery pseudoaneurysm and the duodenum secondary to acute cholecystitis. *Gastrointest Endosc* 2000; 51: 491 – 493
- 2 Lumsden AB, Mattar SG, Allen RC, Bacha EA. Hepatic artery aneurysms: the management of 22 patients. *J Surg Res* 1996; 60: 345 – 350
- 3 Nicholson T, Travis S, Ettles D *et al.* Hepatic artery angiography and embolization for hemobilia following laparoscopic cholecystectomy. *Cardiovasc Intervent Radiol* 1999; 22: 20 – 24

Bibliography

DOI 10.1055/s-0030-1256736

Endoscopy 2011; 43: E323 – E324

© Georg Thieme Verlag KG Stuttgart · New York ·

ISSN 0013-726X

Corresponding author

S. K. Satapathy, MD

Division of Gastroenterology
Hofstra North Shore-LIJ Health system
at Long Island Jewish Medical Center
270-05 76th Avenue
New Hyde Park
New York 11040
USA

Fax: +1-718-343-0128

sanjaya.satapathy@yahoo.com