Human immunodeficiency virus (HIV)-associated duodenal lymphoma

Fig. 1 Endoscopic images of the duodenum showing two ulcerative lesions: **a** second part of the duodenum; **b** inferior duodenal angulus.

A 68-year-old man was referred to our hospital for HIV treatment. He noticed melena on admission. Laboratory data showed anemia, with a hemoglobin level of 8.2 g/dL. The CD4 lymphocyte count was 101 cells/ μ L.

Esophagogastroduodenoscopy (EGD) showed two ulcerative lesions in the second part of the duodenum and in the inferior duodenal angulus (● **Fig. 1**). These well-demarcated lesions consisted of an ulcer with a clean base and a regular elevated margin that had an auricle-like shape and many scattered tiny white spots. Histological examination of the biopsy specimens confirmed the diagnosis of diffuse large B-cell lymphoma (DLBCL). Immunochemical staining was positive for L-26 (CD20) (● **Fig. 2**). Antiretroviral therapy (ART) was started for HIV treatment. After ART, the patient received six cycles of R-CHOP (rituximab, cyclophosphamide, doxorubicin, vincristine, and prednisolone) chemotherapy. Follow-up EGD showed tumor regression. Non-Hodgkin's lymphoma is a common malignancy in HIV-positive individuals,

Fig. 2 Photomicrographs of the biopsy specimen from the duodenal lesion (hematoxylin and eosin stain, $\times 20$): **a** multiple atypical large lymphoid cells infiltrating the duodenal mucosa; **b** large lymphoid cells that are positive for L-26 (CD20).

and the gastrointestinal tract is one of the most common extranodal sites. However, HIV-associated duodenal lymphomas are very rare [1–3]. The endoscopic findings in AIDS-associated lymphoma are reported to be similar to those for gastrointestinal lymphomas in immunocompetent patients: polypoid, bulky lesions or well-defined ulcers are found in combination or as single lesions [4]. The most common HIV-associated lymphomas are Burkitt's lymphoma and DLBCL. Although the incidence of follicular lymphoma in the duodenum is relatively higher than in other portions of the gastrointestinal tract [5], follicular lymphoma has not been reported in HIV-infected patients. Our patient was diagnosed with DLBCL and showed an excellent response

to the R-CHOP chemotherapy administered after ART.

If unique duodenal ulcerative lesions are found, HIV-associated duodenal lymphoma should be considered as a rare differential diagnosis. Furthermore, accurately diagnosing DLBCLs by endoscopic biopsy is very important in such cases because HIV-associated lymphomas of the gastrointestinal tract have a poor prognosis [4] and require aggressive treatment.

Endoscopy_UCTN_Code_CCL_1AB_2AZ_3AB

Competing interests: None

S. Nakazuru¹, T. Yoshio¹, Y. Ogawa²,
K. Yuguchi¹, H. Hasegawa¹, Y. Sakaki-
bara¹, Y. Kodama³, T. Uehira⁴, E. Mita¹

¹ Department of Gastroenterology and
Hepatology, National Hospital Organiza-
tion, Osaka National Hospital, Osaka,
Japan

² Department of Hematology, Osaka City
General Hospital, Osaka, Japan

³ Department of Pathology, National
Hospital Organization, Osaka National
Hospital, Osaka, Japan

⁴ AIDS Medical Center, National Hospital
Organization, Osaka National Hospital,
Osaka, Japan

References

- 1 Bécheur H, Piketty C, Bloch F et al. Endoscopic diagnosis of a duodenocolic fistula due to a non-Hodgkin's lymphoma in a patient with AIDS. *Endoscopy* 1996; 28: 528–529
- 2 Corti M, Villafañe MF, Souto L et al. Burkitt's lymphoma of the duodenum in a patient with AIDS. *Rev Soc Bras Med Trop* 2007; 40: 338–340
- 3 Andhavarapu S, Tolentino AM, Jha C et al. Diffuse large B-cell lymphoma presenting as multiple lymphomatous polyposis of the gastrointestinal tract. *Clin Lymphoma Myeloma* 2008; 8: 179–183
- 4 Heise W, Arastéh K, Mostertz P et al. Malignant gastrointestinal lymphomas in patients with AIDS. *Digestion* 1997; 58: 218–224
- 5 Yoshino T, Miyake K, Ichimura K et al. Increased incidence of follicular lymphoma in the duodenum. *Am J Surg Pathol* 2000; 24: 688–693

Bibliography

DOI 10.1055/s-0030-1256940

Endoscopy 2011; 43: E384–E385

© Georg Thieme Verlag KG Stuttgart · New York ·
ISSN 0013-726X

Corresponding author

S. Nakazuru, MD

Department of Gastroenterology and Hepatology
National Hospital Organization
Osaka National Hospital
2-1-14 Houenzaka
Chuo-ku
Osaka City
Osaka 540-0006
Japan
Fax: +81-6-69463569
nakazuru@onh.go.jp