D.-G. YU, B.-J. LI, S.-F. ZHENG, B.-T. GUAN, B.-Q. WANG, Z.-J. SHI* (PEKING UNIVERSITY, BEIJING, STATE KEY LABORATORY OF ORGANOMETALLIC CHEMISTRY, SHANGHAI AND SICHUAN NORMAL UNIVERSITY, CHENGDU, P. R. OF CHINA)
 Direct Application of Phenolic Salts to Nickel-Catalyzed Cross-Coupling Reactions with Aryl Grignard Reagents
Angew. Chem. Int. Ed. **2010**, 49, 4566–4570.

Nickel-Catalyzed Cross-Coupling of Aryl Grignard Reagents to Phenolate Salts

Selected examples:

Catalytic pathway:

Significance: The first successful cross-coupling of 2-naphthol metal salts with various aryl Grignard reagents has been demonstrated. The process is atom-economical and gives a convenient access to various naphthalene derivatives.

Comment: It is important to note that the halide substituent on the Grignard reagent is critical to the reaction and a bromide was found to be the best. The most efficient solvent system is a mixture of toluene and diisopropyl ether (3:1), most likely because these solvents retain the metallic core framework due to their low coordinating ability.

SYNFACTS Contributors: Paul Knochel, Andreas J. Wagner
 Synfacts 2010, 9, 1051-1051 Published online: 23.08.2010
 DOI: 10.1055/s-0030-1257897; Reg-No.: P09710SF