Z. SUN, K.-W. HUANG, J. WU* (NATIONAL UNIVERSITY OF SINGAPORE, SINGAPORE AND KING ABDULLAH UNIVERSITY OF SCIENCE AND TECHNOLOGY, THUWAL, KINGDOM OF SAUDI ARABIA)

Soluble and Stable Zethrenebis(dicarboximide) and Its Quinone

Org. Lett. **2010**, *12*, 4690-4693.

Soluble and Stable Zethrenebis(carboximide) and Its Quinone

Significance: Despite first being synthesized decades ago by Clar and co-workers, zethrene only recently began attracting major interest due to its potentially interesting optoelectronic properties. Poor air and light sensitivity of unsubstituted zethrene would not have aided its cause. A more convenient route to zethrene was found in the 1960's during attempts to synthesize and isolate the unstable tetrahydrodinaphtho[10]annulene, a precursor of zethrene. This was not successfully done until last year.

Comment: Employing the same strategy discovered by Staab and Sondheimer in the 1960's, the authors take advantage of the automatic transannular cyclization of the tetrahydrodinaphtho[10]annulene to form the butadiene core of the zethrene. Meanwhile, the overall improved stability of **1** no doubt makes it an interesting compound for further optoelectronic studies. Extremely notable as well is that attempts to brominate **1** using *N*-bromosuccinimide in the presence of DMF unexpectedly leads to the formation of the respective quinone **2**.

SYNFACTS Contributors: Timothy M. Swager, Joel Batson
Synfacts 2010, 12, 1357-1357 Published online: 22.11.2010
DOI: 10.1055/s-0030-1258940; Reg-No.: S14210SF