J. M. BAUMES, J. J. GASSENSMITH, J. GIBLIN, J.-J. LEE, A. G. WHITE, W. J. CULLIGAN, W. M. LEEVY, M. KUNO, B. D. SMITH* (UNIVERSITY OF NOTRE DAME, USA)
 Storable, Thermally Activated, Near-Infrared Chemiluminescent Dyes and Dye-Strained Microparticles for Optical Imaging
Nat. Chem. **2010**, 2, 1025-1030.

Supramolecular Glow-Sticks

Significance: This group has previously demonstrated the formation of rotaxanes of far-red (near-infrared) emitting squaraine dyes as a method for creating highly emissive functional materials for biological applications. The emission at ca. 750 nm is in the range where biological tissues have reasonable transparency. The chemiluminescent nature of these dyes and the absence of background can enable imaging with single-photon counting methods.

Comment: The formation of the anthracene peroxide complex and the thermal elimination to produce photons is an elegant method for creating an imaging agent. The ability to functionalize the macrocycle and the squaraine guest can be used to create selective contrast, and there are exciting prospects for biological imaging with the supramolecular devices.

SYNFACTS Contributors: Timothy M. Swager
 Synfacts 2011, 2, 0149-0149 Published online: 19.01.2011
 DOI: 10.1055/s-0030-1259255; Reg-No.: S17510SF

2011 © THIEME STUTTGART • NEW YORK

Category

Synthesis of
Materials and
Unnatural Products

Key words

fluorescence

rotaxanes

host-guest systems

SYNFACTS
of the month

This document was downloaded for personal use only. Unauthorized distribution is strictly prohibited.