SYNLETT Spotlight 345

Iodine Monochloride

Compiled by Luana Silva Magalhães da Forezi

This feature focuses on a reagent chosen by a postgraduate, highlighting the uses and preparation of the reagent in current research

Luana da Silva Magalhães Forezi was born in Recreio/MG, Brazil in 1984. She received her chemistry degree from the Universidade Federal de Juiz de Fora (UFJF), Juiz de Fora/MG, Brazil in 2008. She is currently in the final stages of her M.Sc. studies in organic chemistry, at the Universidade Federal Fluminense, under the supervision of the Professors Maria Cecília Bastos Vieira de Souza and Fernanda da Costa Santos. Her research interests are focused on the synthesis of new compounds, such as ribonucleoside derivatives.

Instituto de Química, Universidade Federal Fluminense, CEP 24020-141 Niterói, Rio de Janeiro, Brazil
E-mail: luaneforezi@hotmail.com

Introduction

Iodine monochloride is a chemical compound with the formula ICl. Because of the difference in the electronegativity of iodine and chlorine, ICl is highly polar and behaves as a source of I^+ . Iodine monochloride is a low melting black or brownish-red solid and widely available (usually in 97–98% purity). It is soluble in alcohol, ether, CS_2 , acetic acid, acetone and pyridine and hydrolyzes in water to HCl and IOH. ICl explodes on contact with potassium metal, mixes with sodium metal, and it can explode if impacted. Its reaction with PCl_3 is extremely exothermic.¹ Iodine monochloride can be easily prepared

by adding an aqueous solution of potassium iodate to potassium iodate dissolved in concentrated HCl, in a closed vessel to avoid the loss of chlorine.² Iodine monochloride is a versatile reagent for the synthesis of a large number of organic compounds being employed, for example, as a source of electrophilic iodine in the synthesis of certain aromatic iodides.³ It cleaves C–Si bonds⁴ and can be used in the electrophilic addition to the double bond in alkenes leading to chloriodoalkanes.⁵ When iodine monochloride is reacted with sodium azide in situ, the iodoazide product is obtained.⁶ Other examples of synthetic applications of this reagent also include electrophilic substitutions in Csp² and electrophilic cyclizations.^{7–9}

Abstracts

(A) Do and Daugulis showed that the iodination of pentachlorobenzene using a combination of iodine chloride and *t*-BuOLi in DMF occurs in excellent yield (90%). Fluorine aromatic compounds like 1,3,5-trifluorobenzene can be either mono- or triiodinated in acceptable yields, depending on the ratio of the halobenzene to the iodine chloride.³

(B) Silylthiophene derivatives as 3,4-difluoro-2,5-bis(trimethylsilyl)thiophene can be submitted to an electrophilic substitution reaction with iodine chloride in anhydrous carbon tetrachloride to give 3,4-difluoro-2,5-diiodothiophene in 80% yield.⁴

(C) The addition of ICl to (*Z*)- and (*E*)-2-butene occurs in an anti-stereospecific manner: the *erythro*-2-chloro-3-iodobutane can be formed by the addition of ICl to (*E*)-1-butene, while the other isomer (*threo*) is formed by the addition of ICl to (*Z*)-1-butene.⁵

(D) In their research efforts to modify citidine nucleosides, Verheyden and Moffatt showed that the treatment of enol ether nucleoside **1** with iodine azide, generated in situ from iodine chloride and sodium azide, afforded the 4'-azido nucleoside derivative **1a**.⁶

(E) Goodman et al. reported the reaction between 2β-carbo(2-fluoroethoxy)-3-{3'-[(*Z*)-2-trimethylstannylethenyl]phenyl}nortropane (**2**) and ICl, leading to compound **4** in good yield. Similar results were obtained in the case of 2β-carbo(2-fluoropropoxy)-3β-{3'-[(*Z*)-2-trimethylstannylethenyl]phenyl}nortropane (**3**).⁷

(F) A set of 4-iodoisocoumarins were efficiently prepared by electrophilic cyclization of *ortho*-(1-alkynyl)benzoates using iodine monochloride in dichloromethane.⁸

(G) Electrophilic cyclization of 2-chalcogene alkynyl anisole using ICl as an electrophile source produced 3-iodo-2-(phenylselenanyl)benzofuran in good yield. Manarim et al. reported that for this type of reaction the product distributions were strongly dependent on the nature of the substituents in the aromatic ring of anisole and on the chalcogen atom directly bonded to the triple bond.⁹

References

- (1) Brisbois, R. G.; Wanke, R. A.; Stubbs, K. A.; Stick, R. V. *Iodine Monochloride*, In *Encyclopedia of Reagents for Organic Synthesis*; John Wiley & Sons: West Sussex, UK, **2004**.
- (2) Beck, M. T.; Ribai, G. *J. Phys. Chem.* **1986**, *90*, 2204.
- (3) Do, H.; Daugulis, O. *Org. Lett.* **2009**, *11*, 421.
- (4) Cardone, A.; Martinelli, C.; Pinto, V.; Babudri, F.; Losurdo, M.; Bruno, G.; Cosma, P.; Naso, F.; Farinola, G. M. *J. Polym. Sci., Part A: Polym. Chem.* **2010**, *48*, 285.
- (5) Schmida, G.; Gordon, J. *Can. J. Chem.* **1986**, *64*, 2171.
- (6) Smith, D. B.; Martin, J. A.; Klumpp, K.; Baker, S. J.; Blomgren, P. A.; Devos, R.; Granycome, C.; Heng, J.; Hobbs, C. J.; Jiang, W.; Laxton, C.; Pogam, S. L.; Leveque, V.; Ma, H.; Maile, G.; Merrett, J. H.; Pichota, A.; Sarma, K.; Smith, M.; Swallow, S.; Symons, J.; Vesey, D.; Najera, I.; Cammack, N. *Bioorg. Med. Chem. Lett.* **2007**, *17*, 2570.
- (7) Stehouwer, J. S.; Jarkas, N.; Zeng, F.; Voll, R. J.; Williams, L.; Camp, V. M.; Malveaux, E. J.; Votaw, J. R.; Howell, L.; Owens, M. J.; Goodman, M. M. *J. Med. Chem.* **2008**, *51*, 7788.
- (8) Roy, S.; Roy, S.; Neuenswamder, B.; Hill, D.; Larock, R. C. *J. Comb. Chem.* **2009**, *11*, 1128.
- (9) Manarim, F.; Roehrs, J. A.; Gay, R. M.; Brandão, R.; Menezes, P. H.; Nogueira, C. W.; Zeni, G. *J. Org. Chem.* **2009**, *74*, 2153.