S. C. MARINESCU, D. S. LEVINE, Y. ZHAO, R. R. SCHROCK,* A. H. HOVEYDA*
 (MASSACHUSETTS INSTITUTE OF TECHNOLOGY, CAMBRIDGE AND BOSTON COLLEGE,
 CHESTNUT HILL, USA)
 Isolation of Pure Disubstituted *E* Olefins through Mo-Catalyzed *Z*-Selective Ethenolysis of Stereoisomeric Mixtures
J. Am. Chem. Soc. **2011**, *133*, 11512–11514.

Purification of Alkene Stereoisomers through *Z*-Selective Ethenolysis

Selected examples:

Two-step synthesis of pure *E*-olefins:

Significance: Many methods exist for preparing disubstituted *E*-olefins, such as the Horner–Wadsworth–Emmons reaction and olefin cross-metathesis. While these reactions have proven to be highly reliable and practical, the undesired *Z*-isomer is often difficult to separate by chromatography or distillation. In the present communication, the authors disclose how pure *E*-olefins can be obtained from an *E/Z* mixture through the selective ethenolysis of the *Z*-isomer.

Comment: The by-products of the reaction are of much lower molecular weight than the starting materials, so the pure *E*-isomer can often be obtained through distillation. The rate of ethenolysis is found to be up to 30 times faster for the *Z*-isomer, explaining why such a highly pure *E*-olefin can be obtained. The scale of the reactions was not mentioned.

SYNFACTS Contributors: Mark Lautens, Stephen G. Newman
 Synfacts 2011, 10, 1083–1083 Published online: 20.09.2011
 DOI: 10.1055/s-0030-1261135; Reg-No.: L11211SF