Does patient history and physical examination predict MRI proven cauda equina syndrome?

Authors Jeremy Fairbank¹, Robin Hashimoto², Andrew Dailey³, Alpesh A Patel³, Joseph R Dettori²

Institutions ¹Nuffield Orthopaedic Centre, Oxford, UK
²Spectrum Research Inc, Tacoma, WA, USA
³University of Utah, Salt Lake City, UT, USA

ABSTRACT

Study design: Systematic review.

Study rationale: While magnetic resonance imaging (MRI) is used as the diagnostic gold standard for cauda equina syndrome (CES), many MRI scans obtained from patients presenting with signs and/or symptoms of CES do not reveal concordant pathology. As a result, the role of the history and physical examination remains unclear when determining which patients require emergent MRI.

Objective or clinical question: Are there elements from the history or physical examination that are associated with CES as established by MRI?

Methods: A systematic review of the literature was undertaken for articles published through April 13, 2011. PubMed, Cochrane, National Guideline Clearinghouse Databases, and bibliographies of key articles were searched. Two independent reviewers reviewed articles. Inclusion and exclusion criteria were set and each article was subject to a predefined quality-rating scheme.

Results: We identified four articles meeting our inclusion criteria. All studies evaluated patients with symptoms suggestive of CES and compared symptoms and/or signs with findings at MRI. The mean prevalence of CES as diagnosed by MRI ranged from 14%–48% of patients. No symptoms or signs reported by more than one study showed high sensitivity and specificity, and all likelihood ratios were low. Symptoms included back/low back pain, bilateral sciatica, bladder retention, bladder incontinence, frequent urination, decreased urinary sensation, and bowel incontinence; signs included saddle numbness and reduced anal tone.

Conclusions: There is low evidence that individual symptoms or signs from the patient history or clinical examination, respectively, can be used to diagnose CES. Additional prospective studies are needed to evaluate whether any single and/or combination of symptoms are associated with a positive diagnosis of CES.

This systematic review was funded by AOSpine.

STUDY RATIONALE AND CONTEXT

Cauda equina syndrome (CES) is often defined by a broad range of symptoms and physical examination findings: back pain, leg pain, weakness, numbness, and bowel or urinary changes. However, many of these symptoms may also be caused by medications, nonspinal pathology, or by spine diseases that can be treated nonemergently.

Cauda equina syndrome is one of the most commonly litigated diagnoses with medicolegal concerns regarding the impact of a missed or delayed diagnosis. Given the lack of clearly defined symptoms and signs as well as the legal risks associated with perceived delays in diagnosis, magnetic resonance imaging (MRI) is frequently obtained in patients presenting with these symptoms.

While MRI, coupled with patient history and examination, remains the diagnostic gold standard, it comes at a high cost with many patients demonstrating no concordant pathology. Given the need to accurately diagnose CES while balancing the cost of diagnosis, the role of patient history and physical examination needs to be clarified.

There is a problem of “true diagnosis” here in that CES is only really established when treatment has failed. At presentation with symptoms and signs of CES, not all subjects have CES (perhaps 40% in terms of MRI). At investigation (usually MR) a judgment has to be made if there is cauda equina decompression that might be relieved by surgery. Surgical success is if there are no residual symptoms, but there cannot be complete certainty that this would not have happened by natural resolution. Surgical failure is CES. We have chosen to use MR as the gold standard since it is at this stage that a surgical solution has to be planned and executed.

CLINICAL QUESTION

Are there elements of the patient history or physical examination that are associated with CES as established during imaging?

METHODS

Study design: Systematic review.

Sampling: PubMed, Cochrane Collaboration Database, bibliographies of key articles. Dates searched: through April 13, 2011.

Inclusion criteria: (1) adults presenting with clinical features suggestive of CES; (2) diagnostic studies evaluating elements from patient history (symptoms) and/or from the physical examination (signs); (3) diagnostic imaging (MRI, CT, or myelography) or findings at surgery as the gold standard.

Exclusion criteria (Fig 1): (1) patients < 18 years old; (2) patients with trauma, inflammatory cause, spondylolisthesis, Paget disease, osteochondrosis, congenital malformation, visceral diseases, or cancer; (3) confirmed CES in all patients; (4) gold standard used is composite of surgery or imaging with index test; and (5) therapeutic studies.

Elements from history and physical examination:

Symptoms

- Back pain
- Leg pain/sciatica
- Neurological symptoms (weakness; bowel or bladder)
- Aggravating factors
- Alleviating factors

Signs

- Neurological examination (motor or sensory)
- Reflexes
- Rectal tone

Outcomes:

- Prevalence of CES according to MRI findings
- Summary of sensitivity, specificity, and likelihood ratios for symptoms and signs reported by more than one study

Analysis: Prevalence, sensitivities, specificities, and likelihood ratios were recorded or calculated from the available data. Overall strength of the evidence was assessed using GRADE criteria.

Additional methodological and technical details are provided in the Web Appendix at www.aospine.org/lebsj

Fig 1 Results of literature search.

RESULTS

- From 77 citations, 8 underwent full-text review. Four [1–4] met the inclusion criteria (**Table 1**): 1 prospective (LoE-III) [2] and 3 retrospective (LoE-IV) studies [1, 3, 4]. (See **Table 2** of the Web Appendix).
- Patients with suspected CES underwent diagnostic MRI (**Table 1**, Web appendix), mean prevalence CES: 14%–48% (**Tables 3–4**, Web Appendix) [1–4].

Symptoms (**Table 3**)

- Back pain and bowel incontinence [1–4] had high sensitivities, low specificities, and low LR_s [1, 3, 4].
- Bilateral sciatica had lower sensitivity, higher specificity, and low LR_s [1–4]
- Bladder incontinence, bladder retention, decreased urinary sensation, and frequent urination had varying sensitivities and specificities and low LR_s [1–4].

Signs (**Table 4**)

- Saddle numbness had varying sensitivities and specificities and low LR_s [2, 3].
- Reduced anal tone had low sensitivity, high specificity, and low LR_s [1–4].
- Additional symptoms and signs reported by one study each are summarized in **Tables 3** and **4** of the Web Appendix.

CLINICAL GUIDELINES

None found.

Table 1 Patient and treatment characteristics of included studies investigating diagnostic accuracy of elements of patient history or physical examination compared with imaging or findings at surgery.*

Author	Study design (LoE)	Demographics	Symptoms/patient history	Signs/physical examination	Reference standard; criteria positive CES diagnosis	Study objective; inclusion/exclusion
Balasubramanian et al [1]; (2010) Domen et al [3]; (2009)	Retrospective diagnostic study	N=80 Male: NR Mean age: NR (range, 21–90 y) (57% in 4th and 5th decades)	– Back pain: 90% (72/80) – Unilateral leg pain: 75% (60/80) – Bilateral leg pain: 11% (8/80) – Bladder incontinence: 48% (33/69) – Bladder retention: 18% (14/79) – Bowel incontinence: 15% (12/79) – Mean duration of symptoms: NR	– Saddle sensory deficit/saddle anesthesia: 26% (20/77) – Reduced anal tone/lax anal sphincter: 9% (7/75)	MRI Details of lumbar MRI scans: – NR Criteria for CES diagnosis (MRI): – Canal compromise > 75%	Objective – To evaluate efficacy of clinical assessment in diagnosing CES Inclusion – All patients seen by on-call spine surgery team presenting with clinical features of CES (Jan–Dec 2010) Exclusion – NR

Table 1 (cont) Patient and treatment characteristics of included studies investigating diagnostic accuracy of elements of patient history or physical examination compared with imaging or findings at surgery.*

Author	Study design (LoE)	Demographics	Symptoms/patient history	Signs/physical examination	Reference standard; criteria positive CES diagnosis	Study objective; inclusion/exclusion
Bell et al [2] (2007); Balasubramanian et al [1] (2010)	Prospective diagnostic study	N=23 Male: 61% Mean age: 39 (range, 17–59) y	<ul style="list-style-type: none"> – Back pain: 100% (23/23) – Sciatica: 74% (17/23) – Unilateral sciatica: 57% (13/23) – Bilateral sciatica: 17% (4/23) – Urinary symptoms (any): 100% (23/23) – Urinary incontinence: 26% (6/23) – Painful urinary retention: 30% (7/23) – Painless urinary retention: 4% (1/23) – Decreased sensation of urination: 30% (7/23) – Frequent urination: 30% (7/23) – Erectile dysfunction: 0% males (0/14) – Mean duration of back pain: 745 (range, 1–4500) d – Mean duration of urinary symptoms: 4 (range, 1–24) d 	– NR	<p>MRI</p> <p>Details of lumbar MRI scans:</p> <ul style="list-style-type: none"> – NR <p>Criteria for CES diagnosis (MRI):</p> <ul style="list-style-type: none"> – Disc prolapse causing significant cauda equina/thecal distortion 	<p>Objective</p> <ul style="list-style-type: none"> – To determine whether any single symptom can be used to accurately predict presence of CES on subsequent MRI <p>Inclusion</p> <ul style="list-style-type: none"> – All patients referred to one neurosurgical center from primary care with suspected diagnosis of CES <p>Exclusion</p> <ul style="list-style-type: none"> – Patients who admitting neurosurgical registrar was confident did not have signs or symptoms of CES; patients who had already received diagnostic imaging
	Retrospective diagnostic study	N=58 Male: NR Mean age: NR	<ul style="list-style-type: none"> – Low back pain <1 mo: 74% (43/58) – Bilateral sciatica: 33% (19/58) – Bladder incontinence: 40% (23/58) – Bladder retention: 64% (37/58) – Decreased sensation of urination: 36% (21/58) – Frequent urination: 10% (6/58) – Bowel incontinence: 16% (9/58) – Mean duration of symptoms: NR 	<ul style="list-style-type: none"> – Urinary retention > 500 mL: 21% (8/39 tested) – Saddle anesthesia/numbness: 36% (21/58) – Decreased anal sphincter tone: 16% (9/58) – Decreased anal sphincter reflex: 40% (23/58) – Lasègè straight leg test positive: 48% (28/58) – Loss of motor function in leg: 50% (29/58) – Loss of motor function in foot raisers: 33% (19/58) – Decreased ankle reflex: 29% (17/58) 	<p>MRI</p> <p>Details of lumbar MRI scans:</p> <ul style="list-style-type: none"> – All scans done on same 1.5-T clinical MRI scanner (Intera; Phillips, Best, the Netherlands) using following sequences (done according to in-house lumbar protocol): sagittal T1-weighted MRI (12 slides; SL 4.00 mm; TR 550 ms; TE 13 ms), sagittal T2-weighted MRI (12 slides; SL 4.0 mm; TR 3191 ms; TE 130 ms), axial driven-equilibrium (T2) MRI (50 slides; SL 2.0 mm; TR 700 ms; TE 1010 ms) and MR cauda equina myelography <p>Criteria for CES diagnosis (MRI):</p> <ul style="list-style-type: none"> – Presence of cauda equina compression 	<p>Objective</p> <ul style="list-style-type: none"> – To determine whether the presence or absence of CES can be accurately predicted by certain clinical characteristics in the hopes of avoiding unnecessary urgent MRI scans <p>Inclusion</p> <ul style="list-style-type: none"> – All patients who underwent urgent MRI imaging for suspected CES at hospital neurology department (Jan 2003–Dec 2007) <p>Exclusion</p> <ul style="list-style-type: none"> – Patients with pre-medical history of recent or extensive malignant disease; patients with metastatic disease as cause of CES

Table 1 (cont) Patient and treatment characteristics of included studies investigating diagnostic accuracy of elements of patient history or physical examination compared with imaging or findings at surgery.*

Author	Study design (LoE)	Demographics	Symptoms/ patient history	Signs/physical examination	Reference standard; criteria positive CES diagnosis	Study objective; inclusion/exclusion
Rooney et al [4] (2009); Domen et al [3] (2009)	Retrospective diagnostic study	N=98† Male: 27% (18/66)† Mean age: 43 y	– Low back pain: 94% (58/62)	– Saddle numbness: 55% (34/66)	MRI	Objective
			– Sciatica: 89% (49/55)	– Leg numbness: 80% (49/61)	Details of lumbar MRI scans: – NR	– To assess whether any simple clinical characteristics are able to distinguish patients with CES from those without
			– Unilateral sciatica: 60% (33/55)	– Loss of power (unilateral or bilateral): 59% (38/64)	Criteria for CES diagnosis (MRI):	Inclusion
			– Bilateral sciatica: 29% (16/55)	– Unilateral loss of power: 33% (21/64)	– Relevant abnormality on scan (details NR)	– All patients referred to neurosurgical department at hospital with symptoms suggestive enough of CES to warrant MRI (Mar– Dec 2004)
			– Leg numbness: 83% (33/40)	– Bilateral loss of power: 19% (12/64)		
			– Unilateral leg numbness: 53% (21/40)	– Loss of reflexes (unilateral or bilateral): 41% (26/64)		Exclusion
			– Bilateral leg numbness: 30% (12/40)	– Unilateral loss of reflexes: 22% (14/64)		– Readmissions of same patient; previous MRI scan in different hospital; on clinical reevaluation CES seemed unlikely; incomplete records (32 patients had incomplete records)
			– Bladder incontinence: 56% (27/48)	– Bilateral loss of reflexes: 19% (12/64)		
			– Bladder retention: 43% (17/40)	– Increase in reflexes (unilateral or bilateral): 14% (9/64)		
			– Decreased sensation of urination: 43% (18/42)	– Unilateral increase in reflexes: 2% (1/64)		
			– Bowel incontinence: 8% (4/48)	– Bilateral increase in reflexes: 13% (8/64)		
			– Bowel retention: 13% (5/40)			
			– Decreased fecal sensation: 5% (2/42)			
			– Mean duration of symptoms: NR			

* CES indicates cauda equina syndrome; NR, not reported; and MRI, magnetic resonance imaging.

† Complete records available for 66 of 98 patients, the remaining 22 were excluded; all data reported for n = 66 patients with complete records.

ILLUSTRATIVE CASE

A 37-year-old woman had acute onset of low back pain while lifting at work. During 4 days she experienced pain down her legs, bilateral foot weakness, and perineal numbness. She could no longer sense when her bladder was full and began to wear a diaper. On presentation, strength testing revealed plantar flexion, dorsiflexion, and extensor hallucis longus 0/5 on the right, dorsiflexion and hallucis longus (EHL) 4/5 and plantar flexion 5/5 on the left. She had saddle anesthesia, diminished rectal tone, and more than 1000 cc of retained urine upon placing a urinary catheter.

Emergent MRI revealed a massive L4–5 disc herniation causing cauda equina compression (**Fig 2**). Despite the delay in referral, she was taken urgently to the operating room for laminectomy and disc excision. She made a full-motor recovery over 2 years but she still has sacral numbness and manually expresses her bladder, although she no longer needed a catheter.

Fig 2 **a** Axial (left); **b** sagittal (right). T2 magnetic resonance imaging reveals a massive disc extrusion from the L4–5 level associated with signs and symptoms of cauda equina syndrome.

DISCUSSION

Limitations

- Small number of poor studies available (three studies are level IV; one level III).
- No indication in any study whether the reference test (MRI) was interpreted in a manner that was blind to the results of the patient history and physical examination.
- None of the studies provided sufficient details for replication of both the tests and MRI.
- None of the symptoms or signs reported had a likelihood ratio with a magnitude that would suggest ruling in or out CES given the pretest probability of disease.
- The literature did not define objective, reliable clinical criteria for the diagnosis of CES.
- There are insufficient data in the literature to determine the relationship between signs and symptoms of CES and the timing or severity (ie, complete vs incomplete) of the disease.
- While MRI is considered the standard in the diagnosis of CES, the literature reveals a lack of an objective, reliable system of classifying MRI findings. Furthermore, the subjective MRI findings reported in the literature may not be specific to CES.
- Insufficient data is present in the literature to identify signs or symptoms that correlated with either negative or positive MRI studies.
- Prospective studies evaluating larger cohorts of patients are needed to more definitively determine whether any individual or combination of signs and/or symptoms is associated with CES.
- Prospective studies should comprise objective, reliable definitions for CES, including timing and severity of disease, as well as a reliable classification of MRI findings.

EVIDENCE SUMMARY

Question 1: Are there elements of the patient history that are associated with cauda equina syndrome (CES) as established during imaging?		
Outcomes	Strength of evidence	Conclusions/comments
Sensitivity, specificity, and likelihood ratio	<p>Further research is likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.</p>	One prospective and three retrospective cohort studies were identified that compared symptoms as established during patient history taking to MRI findings regarding CES. All symptoms identified had low likelihood ratios. No studies evaluated a combination of symptoms.
Question 2: Are there elements of the physical examination that are associated with CES as established during imaging?		
Outcomes	Strength of evidence	Conclusions/comments
Sensitivity, specificity, and likelihood ratio	<p>Further research is likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.</p>	One prospective and three retrospective cohort studies were found that compared signs as established during the physical examination to the CES diagnosis according to MRI. All symptoms identified had low likelihood ratios. No studies evaluated a combination of signs.

REFERENCES

1. **Balasubramanian K, Kalsi P, Greenough CG, et al** (2010) Reliability of clinical assessment in diagnosing cauda equina syndrome. *Br J Neurosurg*; 24(4):383–386.
2. **Bell DA, Collie D, Statham PF** (2007) Cauda equina syndrome: what is the correlation between clinical assessment and MRI scanning? *Br J Neurosurg*; 21(2):201–203.
3. **Domen PM, Hofman PA, van Santbrink H, et al** (2009) Predictive value of clinical characteristics in patients with suspected cauda equina syndrome. *Eur J Neurol*; 16(3):416–419.
4. **Rooney A, Statham PF, Stone J** (2009) Cauda equina syndrome with normal MR imaging. *J Neurol*; 256(5):721–725.

- CES is a description of a neural deficit best to be avoided in the first place as it describes an ongoing neurological deficit.
- CES can be defined as the sudden loss of function of the lumbar and lumbosacral plexus below the conus medullaris due to a number of conditions, such as massive disc herniation, tumor, trauma, and by other forms of neural canal compromise.
- The traditional findings of acute loss of bowel and bladder control function, sexual dysfunction, lower extremity weakness, and numbness in the perianal region and lower extremities describe an active process of neurological damage underway for which recovery is unpredictable. Recognition of an immanent CES is clearly preferable to waiting for actual manifestations of CES to occur.
- CES remains a major challenge regarding best pathway to timely diagnosis due to a plethora of presentations and large number of confounding factors.
- Lumbar MRI remains the preferred diagnostic modality to identify neural space compromise.

In the era of attempting to develop pathways for use of lumbar neural imaging for patients with low back pain based on their clinical presentations, the possibility of missing an evolving CES remains a real challenge. A review of the physical examination findings reported in this systematic review by Fairbanks et al clearly show the main constant of physical findings with patients who ended up experiencing CES was 'low back pain. A smaller number of patients presented with bilateral leg pain. All other physical examination findings pretty much described an ongoing evolving CES—the condition all agreed that should be avoided in the first place. The authors clearly found that there was a dearth of reliable patient complaints or physical examination findings in the literature to use as a 'red flag' in the development of diagnostic pathways for low back pain. There also appears to be poor correlation of the amount of canal compromise and the evolution as well as severity of neurological deficits.

In conclusion, we can all agree on some points:

- CES as a worst-case scenario for a number of conditions, such as disc herniation, clearly begs for further prospective investigation with quality-data collection and analysis.
- The neurophysiological basis of CES would benefit from further basic science study.
- Low back pain continues to present a major challenge to simplified diagnostic and care pathways due to its highly nonspecific nature.

EBSJ thanks Fairbank and colleagues for tackling this controversial subject and providing a welcome foundation for further investigation.

EDITORIAL PERSPECTIVE

The subject of cauda equina syndrome (CES) continues to provide ample controversy to all involved in care of spine patients, scientific study, and education. This includes the responses of our reviewers who were split about the best way to approach the subject and then interpret the available literature and what conclusions to draw from the findings. All agree on the following: