Risk adjustment – never enough or too much?

J. Nicholl

Qualitätsmanagement

Keywords

- ▶ risk adjustment
- ▶ hospital comparisons
- ▶ constant risk fallacy

Schlüsselwörter

- ▶ Risikoadjustierung
- ▶ Krankenhausvergleich
- ▶ constant risk fallacy

Comparisons of health outcomes between people, services, institutions, and populations are used to assess the relative quality of health care. However, these non-experimental comparisons may be confounded by differences in casemix factors associated with the outcome. Casemix adjustment is used to create a fair comparison – but does it? Simulation experiments suggest that casemix adjustment can make the bias worse in non-experimental studies. How is this possible? This talk will explore the problem that casemix variables may have a different effect in different populations – this is the constant risk fallacy. Examples of non-constant risk in hospital comparisons will be presented and possible solutions will be explored.

Institut

School of health
and Related research,
University of Sheffield,
England

Bibliografie

DOI 10.1055/s-0031-1286095
Dtsch Med Wochenschr 2011;
136: S66 · © Georg Thieme Ver-
lag KG Stuttgart · New York ·
ISSN 0012-0472

Korrespondenz

Prof. Jon Nicholl, DSc
Dean, School of health and
Related research
University of Sheffield
Western Bank
Sheffield S10 2TN
England