SYNLETT Spotlight 372

Sodium Sulfide

Compiled by Fei-Ping Ma

This feature focuses on a reagent chosen by a postgraduate, highlighting the uses and preparation of the reagent in current research

Fei-Ping Ma was born in 1988 in Shijiazhuang, Hebei Province, P. R. of China. She graduated from the Huihua College of Hebei Normal University in 2010 and received her B.Sc. degree in chemistry. Presently she is working as a postgraduate towards her M.Sc. under the supervision of Professor Zhan-Hui Zhang at the Hebei Normal University. Her research interest focuses on the development of new synthetic methodologies of heterocyclic compounds.

The College of Chemistry & Material Science, Hebei Normal University, Shijiazhuang 050016, P. R. of China
E-mail: mafp0312@163.com

Introduction

Sodium sulfide is pink to yellowish solid with a rotten egg-like odor. It is readily soluble in water, slightly soluble in alcohol and insoluble in ether. Anhydrous sodium sulfide can ignite spontaneously when exposed to air.¹ It is used in chemical manufacturing as a sulfonation and sulfomethylation agent. It is used in the production of rubber chemicals, sulfur dyes, and other chemical compounds. It has also been used as a reducer for the reduction of nitro compounds to the corresponding amines.² It was found to be useful for the conversion of carboxylic acids into thioacids³ and alkenoyl ketene dithioacetals into the corresponding [5+1]-annulation products⁴ as well as for the synthesis of thiofuranose,⁵ cyclic dithiocarbonates,⁶

tetrahydrothiophene derivatives,⁷ 4-bromo-2-nitrobenzenethiol,⁸ α-lipoic acid,⁹ thieno[2,3-*c*]pyrazoles,¹⁰ unsaturated thiacycrown ethers,¹¹ thieno[3,2-*c*]cinnoline,¹² monomeric cyclic diketosulfides,¹³ thiophenes,¹⁴ 2*H*-thiopyran compounds,¹⁵ 4*H*-thiochromen-4-ones¹⁶ and benzo[*b*]thiophenes.¹⁷ It can also act as an atom-economical inorganic nucleophile in transition-metal-catalyzed allylation substitutions.¹⁸ Willgerodt–Kindler reaction between anilines and benzaldehydes has also been achieved by the use of Na₂S·9H₂O as a base catalyst.¹⁹

Sodium sulfide is commercially available and it can be readily prepared by reduction of Na₂SO₄ with carbon or through a solid–gas reaction of a sulfidizing gas mixture of COS, CS₂, and S₂ with the reactant of Na₂CO₃.²⁰

Abstracts

(A) Thionation of Carbonyl Compounds:

Salama and co-workers have found that a combination of tetrachlorosilane and sodium sulfide in acetonitrile is an efficient thionating reagent for aromatic aldehydes to afford the corresponding thioaldehydes as trimers in good yields in the absence of catalysis. Under these mild conditions α,β-unsaturated ketones reacted with SiCl₄/Na₂S in the presence of a catalytic amount of CoCl₂·6H₂O to give the respective disulfides via β-mercapto ketones.²¹

(B) Regioselective Ring Opening of Epoxides:

Na₂S can be applied as a mild sulfur nucleophile for regioselective ring opening of epoxides to give the corresponding bis[β-hydroxyalkyl]sulfides in the presence of poly[*N*-(2-aminoethyl)acrylamido]trimethyl ammonium chloride resin as a phase-transfer catalyst.²² The reaction of 2,2-bis(trifluoromethyl)oxiranes with aqueous solution of Na₂S leads to the formation of S[CH₂C(CF₃)₂OH]₂.²³

(C) Divinylsulfides:

A convenient and practical method for the direct synthesis of bis(arylvinyl)sulfides by the addition of sodium sulfide to arylacetylenes has been developed. These sulfides can be converted chemoselectively into the sulfoxides and sulfones.²⁴

(D) 2-Trifluoromethyl Benzothiophenes:

Li et al. reported a practical protocol for the selective synthesis of 2-trifluoromethyl benzothiophenes by copper-catalyzed thiolation annulation reaction of 1,4-dihalides with Na_2S . This protocol allows the formation of two C–S bonds in a one pot reaction through the thiolation annulation of various 1,4-dihalides, including less active dichlorides.²⁵

(E) Substituted 2,3,6,7-Tetrahydrothiopyrano[2,3-b]thiopyran-4,5-diones:

Pan et al. reported a one-pot approach for the synthesis of disubstituted bicyclic thia-heterocycles via formal double [5+1] annulation of dialkenylketene dithioacetals using sodium sulfide nonahydrate in DMF.²⁶

(F) Symmetrical Alkyl Disulfides:

A mild and practical method for the synthesis of symmetrical alkyl disulfides from alkyl halides and a mixture of Na_2S with sulfur using didecyltrimethylammonium bromide (DDAB) as a phase-transfer catalyst has been developed.²⁷ The reactivity order was found to be: benzyl halide > primary alkyl halide > secondary alkyl halide > tertiary alkyl halide > aryl halide.

(G) Angular Anthrathiophenediones:

In addition to the above cases, the cyclocondensation of available vic-alkynylchloroanthraquinones with Na_2S is a convenient method for the synthesis of angular anthrathiophenediones.²⁸

References

- (1) Young, J. A. *J. Chem. Edu.* **2009**, 86, 919.
- (2) (a) Akue-Gedu, R.; Gautret, P.; Lelieur, J.-P.; Rigo, B. *Synthesis* **2007**, 3319. (b) Zimmermann, V.; Müller, R.; Bräse, S. *Synlett* **2008**, 278.
- (3) El-Faham, A.; Khattab, S. N. *Synlett* **2009**, 886.
- (4) Pan, L.; Liu, Q. *Synlett* **2011**, 1073.
- (5) Qing, F.-L.; Zheng, F. *Synlett* **2011**, 1052.
- (6) Crivillers, N.; Oxtoby, N. S.; Mas-Torrent, M.; Veciana, J.; Rovira, C. *Synthesis* **2007**, 1621.
- (7) Periasamy, M.; Ramani, G.; Muthukumaragopal, G. P. *Synthesis* **2009**, 1739.
- (8) Pirat, C.; Ultré, V.; Lebegue, N.; Berthelot, P.; Yous, S.; Carato, P. *Synthesis* **2011**, 480.
- (9) Kaku, H.; Okamoto, N.; Nishii, T.; Horikawa, M.; Tsunoda, T. *Synthesis* **2010**, 2931.
- (10) Eller, G. A.; Vilkauskaitė, G.; Arbačiauskienė, E.; Sačkus, A.; Holzer, W. *Synth. Commun.* **2011**, 41, 541.
- (11) Sun, D.-Q.; Yang, J.-K. *Synthesis* **2011**, 2454.
- (12) Vinogradova, O. V.; Sorokoumov, V. N.; Balova, I. A. *Tetrahedron Lett.* **2009**, 50, 6358.
- (13) Miyahara, Y. *J. Org. Chem.* **2006**, 71, 6516.
- (14) Thomae, D.; Kirsch, G.; Seck, P. *Synthesis* **2007**, 1027.
- (15) Banerji, A.; Biswas, P. K.; Bandyopadhyay, D.; Gupta, M.; Prangé, T.; Neuman, A. *Phosphorus, Sulfur Silicon Relat. Elem.* **2009**, 184, 3199.
- (16) Willy, B.; Müller, T. J. J. *Synlett* **2009**, 1255.
- (17) Saito, M.; Yamamoto, T.; Osaka, I.; Miyazaki, E.; Takimiya, K.; Kuwabara, H.; Ikeda, M. *Tetrahedron Lett.* **2010**, 51, 5277.
- (18) Zheng, S. C.; Huang, W. Q.; Gao, N.; Cui, R. M.; Zhang, M.; Zhao, X. M. *Chem. Commun.* **2011**, 47, 6969.
- (19) Okamoto, K.; Yamamoto, T.; Kanbara, T. *Synlett* **2007**, 2687.
- (20) Güler, H.; Kurtuluş, F.; Kadan, İ.; Morkan, A.; Akin, S. *Phosphorus, Sulfur Silicon Relat. Elem.* **2006**, 181, 1371.
- (21) Salama, T. A.; El-Ahl, A.-A. S.; Elmorsy, S. S.; Khalil, A.-G. M.; Ismail, M. A. *Tetrahedron Lett.* **2009**, 50, 5933.
- (22) Mahdavi, H.; Tamami, B. *Phosphorus, Sulfur Silicon Relat. Elem.* **2005**, 180, 1929.
- (23) Petrov, V. A.; Marshall, W. J. *Fluorine Chem.* **2011**, 132, 41.
- (24) Paradies, J. *Synthesis* **2010**, 947.
- (25) Li, C.-L.; Zhang, X.-G.; Tang, R.-Y.; Zhong, P.; Li, J.-H. *J. Org. Chem.* **2010**, 75, 7037.
- (26) Pan, W.; Dong, D.; Ouyang, Y.; Wu, R.; Yang, Y.; Liu, Q. *Synthesis* **2007**, 2115.
- (27) Sonavane, S. U.; Chidambaram, M.; Almog, J.; Sasson, Y. *Tetrahedron Lett.* **2007**, 48, 6048.
- (28) Ivanchikova, I. D.; Lebedeva, N. I.; Shvartsberg, M. S. *Synthesis* **2004**, 2131.