Y. NISHIMOTO, M. TAKEUCHI, M. YASUDA, A. BABA* (OSAKA UNIVERSITY, SUITA, JAPAN)

Regio- and Stereoselective Carbobismuthination of Alkynes

Angew. Chem. Int. Ed. **2012**, 51, 1051–1054.

anti-Carbobismuthination of Alkynes

R¹ = Ph, 1,3-di-*t*-BuC₆H₃, 4-ClC₆H₄, Hex

R²/R³ = H, Me, Ph

electrophile = I₂, TsCl, PhSPh, RCOCl

Selected examples:

Significance: The first carbobismuthination of alkynes has been accomplished by the reaction of an alkyne, BiBr₃, and a ketene silyl acetal to produce an alkenylbismuth compound with high stereo- and regioselectivity. The Br₂Bi group in the alkenylbismuth compounds can be substituted by I, Ts and SPh groups, and palladium-catalyzed cross-couplings with acid chlorides have been performed successfully.

Comment: The reaction of BiBr₃ with a phenyl-acetylene derivative and a ketene silyl acetal gives monoalkenylbismuth dibromide as a white solid. X-ray crystallographic analysis of this product reveals that the carbobismuthination takes place regio- and stereoselectively in an *anti*-addition manner.

SYNFACTS Contributors: Paul Knochel, Andreas K. Steib
Synfacts 2012, 8(4), 0427 Published online: 20.03.2012
DOI: 10.1055/s-0031-1290538; Reg-No.: P02112SF

2012 © THIEME STUTTGART • NEW YORK

Category

Metal-Mediated
Synthesis

Key words

bismuth

alkynes

carbobismuth-
ination

SYNFACT
of the month