Thoracic spine osteophyte causing dysphagia

Fig. 1 Polypoid lesion in the esophagus.

Fig. 2 Endoscopic ultrasonography showing erosion of posterior esophageal wall by vertebra (arrow).

Fig. 3 Contrast-enhanced computed tomography showing erosion into the posterior esophageal wall by vertebra (arrow).

Fig. 4 Contrast-enhanced computed tomography with positive oral contrast in esophagus could not demonstrate the lesion in the posterior esophageal wall.

A 65-year-old man, a chronic smoker and alcoholic with a diagnosed adenocarcinoma of the antrum of the stomach, was being further investigated because of dysphagia; endoscopic examination at the referring center showed a polypoid lesion in the mid esophagus. Positron emission tomography (PET) revealed increased uptake of fluorodeoxyglucose in this esophageal lesion. The possibility of a metastasis to the esophagus from the gastric malignancy was considered. However, biopsy from the esophageal lesion revealed features of chronic inflammation. A repeat gastroscopy was done and a polypoid lesion was observed in the mid esophagus (● Fig. 1). Histopathological examination of the biopsy specimen from this lesion

again revealed features of chronic inflammation. Contrast-enhanced computed tomography (CECT) of the chest with intravenous and a positive oral contrast revealed a dilated esophagus. Radial endoscopic ultrasound examination of the esophagus revealed that the vertebral column was eroding into the posterior esophageal wall at the site of the lesion noted on endoscopy (● Fig. 2). A repeat chest CECT, this time without oral contrast, showed that an anterior osteophyte from the thoracic vertebra was eroding into the esophagus (● Fig. 3). It had not been possible to diagnose it in the previous CECT as during that procedure positive oral contrast was given which obscured the vertebral erosion into the esophagus

(● Fig. 4). A barium esophagogram also documented indentation of the posterior wall of the esophagus by a thoracic vertebra (● Fig. 5).

Anterior osteophytes can occasionally impinge on the anteriorly located esophagus and can cause dysphagia [1–4]. This commonly involves the hypopharynx or the cervical esophagus [1–4]. Involvement of the thoracic esophagus is very rare because the thoracic esophagus is a relatively mobile structure in the posterior mediastinum that can be displaced without being compressed [5].

Endoscopy_UCTN_Code_CCL_1AB_2AC_3AH

Competing interests: None

Fig. 5 Barium esophagogram showing posterior indentation of the esophagus by thoracic vertebra (arrow).

**S. S. Rana¹, D. K. Bhasin¹, C. Rao¹,
R. Gupta², B. Nagi¹, K. Singh¹**

¹ Department of Gastroenterology, Postgraduate Institute of Medical Education and Research, Chandigarh, India

² Department of Surgery, Postgraduate Institute of Medical Education and Research, Chandigarh, India

References

- 1 Hilding DA, Tachdjian MO. Dysphagia and hypertrophic spurring of the cervical spine. *N Engl J Med* 1960; 263: 11–14
- 2 Resnick D, Shaul SR, Robins JM. Diffuse idiopathic skeletal hyperostosis (DISH): Forestier's disease with extraspinal manifestations. *Radiology* 1975; 115: 513–524
- 3 Lin HW, Quesnel AM, Holman AS et al. Hypertrophic anterior cervical osteophytes causing dysphagia and airway obstruction. *Ann Otol Rhinol Laryngol* 2009; 118: 703–707
- 4 Seidler TO, Pérez Alvarez JC, Wonneberger K et al. Dysphagia caused by ventral osteophytes of the cervical spine: clinical and radiographic findings. *Eur Arch Otorhinolaryngol* 2009; 266: 285–291
- 5 Underberg-Davis S, Levine MS. Giant thoracic osteophyte causing esophageal food impaction. *AJR Am J Roentgenol* 1991; 157: 319–320

Bibliography

DOI <http://dx.doi.org/10.1055/s-0031-1291501>
Endoscopy 2012; 44: E19–E20
© Georg Thieme Verlag KG
Stuttgart · New York
ISSN 0013-726X

Corresponding author

S. S. Rana, MD
Department of Gastroenterology
Postgraduate Institute of Medical Education and Research (PGIMER)
Chandigarh
160012 India
Fax: 91-172-2744401
drsurinderrana@yahoo.co.in
sonalisurinder@yahoo.co.in