Signet-ring cell carcinoma of the nonampullary duodenum and proximal jejunum: a case report with an immunohistochemical study

Fig. 1 Endoscopy shows an ulcerated tumor (arrow) of the third portion of the duodenum.

Fewer than 20 cases of signet-ring cell carcinoma (SRCC) of the ampulla of Vater have been reported [1–4]. There have been no reports to date of SRCC of the nonampullary duodenum and ileum. A 71-year-old man presented with nausea. Upper gastrointestinal endoscopy showed two ulcerated tumors (● **Fig. 1**) of the third portion of the duodenum and in the proximal jejunum. Biopsies were taken. The stomach was free from tumor. CT demonstrated brain metastases. The patient was treated by chemoradiotherapy, but died of carcinomatosis 4 months after initial presentation. Histologic examination of the biopsies of both tumors showed apparent SRCC

(● **Fig. 2**). Approximately 80% of the tumor was composed of SRCC cells. Histochemically, the SRCC cells were positive for mucins. Immunohistochemically [5], the SRCC cells were positive for cytokeratin AE1/3, cytokeratin CAM5.2, CK34BE12, cytokeratin 7 (CK7), cytokeratin 8 (CK8), cytokeratin 18 (CK18), cytokeratin 19 (CK19), epithelial membrane antigen (EMA), p53, Ki-67 (labeling = 70%), mucin 1 (MUC1), and mucin 6 (MUC6). They were negative for cytokeratin 5/6 (CK5/6), cytokeratin 14 (CK14), cytokeratin 20 (CK20), carcinoembryonic antigen (CEA), carbohydrate antigen 19–9 (CA19–9), vimentin, p63, thyroid transcription factor 1 (TTF-1), homeobox protein CDX-2 (CDX2), mucin 2 (MUC2), and mucin 5AC (MUC5AC).

The present case is the first reported one of SRCC of the nonampullary duodenum and proximal duodenum. Although the primary site is unclear, the present author thinks that the duodenal tumor was primary and the jejunal tumor was a metastasis. The cytokeratin profile was compatible with pancreatobiliary cytokeratins, which are CK7, CK8, CK18, and CK19 [6]. However, the MUC apomucin profile was not of the pancreatobiliary type [6]. The positive CK34BE12 staining suggests that this high-molecular cytokeratin emerged during carcinogenesis. CK5/6, CK14, and CK20 are known to be absent in the duodenal epithelium. The positive p53 stain-

ing in the present case suggests p53 mutations. The high Ki-67 labeling indicates a high proliferative fraction. The EMA is known to be positive in duodenal epithelium. The negative immunoreactivities of CEA and CA19–9 indicate that this SRCC does produce these cancer-related antigens. TTF-1 and CDX2 indicate that the present tumor is not associated with lung and colon phenotypes.

Endoscopy_UCTN_Code_CCL_1AB_2AZ_3AB

Competing interests: None

T. Terada

Department of Pathology, Shizuoka City Shimizu Hospital, Shizuoka, Japan

References

- 1 Gardner HA, Matthews J, Ciano PS. A signet-ring cell carcinoma of the ampulla of Vater. *Arch Pathol Lab Med* 1990; 114: 1071–1072
- 2 Paplomata E, Wilfong L. Signet ring cell carcinoma of the ampulla of Vater with leptomeningeal metastases: a case report. *J Clin Oncol* 2011; 29 e627–629
- 3 Gheza F, Cervi E, Pulcini G et al. Signet ring cell carcinoma of the ampulla of Vater: demonstration of pancreaticobiliary origin. *Pancreas* 2011; 40: 791–793
- 4 Tas A, Ozer E, Koklu S et al. Signet ring cell carcinoma of the ampulla of Vater: a rare case of acute pancreatitis. *Scand J Gastroenterol* 2011; 46: 126–127
- 5 Terada T, Kawaguchi M, Furukawa K et al. Minute mixed ductal-endocrine carcinoma of the pancreas with predominant intraductal growth. *Pathol Int* 2002; 52: 740–746
- 6 Hruban RH, Kloppel G, Boffetta P et al. Ductal adenocarcinoma of the pancreas. In: Bosman FT, Carneiro F, Hruban RH, Theise ND, eds. *WHO classification of tumours of the digestive system*. Lyon: IARC; 2010: 281–291

Bibliography

DOI <http://dx.doi.org/10.1055/s-0031-1291528>
Endoscopy 2014; 46: E348
© Georg Thieme Verlag KG
Stuttgart · New York
ISSN 0013-726X

Corresponding author

T. Terada, MD, PhD
Department of Pathology
Shizuoka City Shimizu Hospital
Miyakami 1231 Shimizu-Ku
Shizuoka 424-8636
Japan
Fax: +81-54-3341173
piyo0111jp@yahoo.co.jp

Fig. 2 Histologic section of biopsy specimen. Signet-ring cell carcinoma is apparent (H&E, ×200).