A rare case of small bowel intussusception

A 58-year-old man was referred to our institution with a 2-month history of worsening colicky upper abdominal pain associated with eating and relieved by vomiting, and weight loss.

Gastroscopy revealed grade B reflux esophagitis, erosive gastritis, and duodenitis. A *Campylobacter*-like organism (CLO) test was negative for *Helicobacter pylori*, and duodenal biopsies were normal. Empirical triple therapy with amoxicillin, clarithromycin, and omeprazole was administered. An abdominal ultrasound was normal. A barium follow-through showed an unusual jejunal appearance with dilatation also noted (● Fig. 1). Subsequent CT enterography suggested proximal jejunal tethering. Double-balloon enteroscopy (DBE) revealed, 10 cm beyond the ligament of Treitz, a large sessile, ulcerated 3–4 cm

lesion, hemicircumferentially involving the small bowel wall (● Fig. 2). Multiple biopsies were taken and a tattoo was placed proximal to the lesion. Histopathological analysis showed a dense lymphoid infiltrate including aggregates/germinal centers with perivascular activity, highly suggestive of lymphoma. Immunohistochemistry confirmed low grade B-cell mucosa-associated lymphoid tissue (MALT lymphoma).

At 2 weeks after a first cycle with R-CVP (rituximab, cyclophosphamide, vincristine, prednisolone), the patient presented with severe abdominal pain. A computed tomography (CT) scan revealed jejunal intussusception (● Fig. 3), but surgical review recommended conservative management. A second DBE showed no change in the lesion (● Video 1) after three further cycles of R-CVP. For this rea-

son, the patient's chemotherapy was changed to a more aggressive scheme with R-CHOP (rituximab, cyclophosphamide, hydroxydaunorubicin, oncovin, and prednisone) for a further three cycles, and a third DBE revealed a marked improvement with significant regression of the mass (● Fig. 4, ● Video 2).

MALT lymphoma is the commonest extranodal small B-cell non-Hodgkin's lymphoma [1]. Of extranodal lymphomas, 37% occur in the gastrointestinal tract, with 7.5% involving the small bowel [2]. While almost 90% of gastric MALT lymphomas are caused by *Helicobacter pylori* [3], the etiology of jejunal MALT lymphoma is uncertain [4]. DBE is a useful tool for the diagnosis and follow-up of small bowel lymphoma [5].

Fig. 1 Unusual jejunal appearance with dilatation, showed by a barium follow-through (typical coiled spring appearance: large arrow, intussusceptum; small arrow: intussusciens).

Fig. 3 Jejunal intussusception revealed by computed tomography (CT) scan: **a** axial view; **b** coronal view.

Video 1

Double-balloon enteroscopy (DBE) showed no change in the lesion after four cycles of R-CVP (rituximab, cyclophosphamide, vincristine, prednisolone).

Video 2

After three cycles of R-CHOP (rituximab, cyclophosphamide, hydroxydaunorubicin, oncovin, and prednisone), double-balloon enteroscopy (DBE) revealed a marked improvement with significant regression of the mass.

Fig. 2 **a, b** A large sessile, ulcerated 3–4 cm lesion, hemicircumferentially involving the small bowel wall.

Fig. 4 Marked improvement with significant regression of the jejunal lesion at double-balloon enteroscopy (DBE).

Endoscopy_UCTN_Code_CCL_1AC_2AC

Competing interests: The first author has a Research Grant in deep enteroscopy released by Imotec/Fujinon.

A. Murino¹, E. J. Despott¹, A. Hansmann², P. Heath³, C. Fraser¹

¹ Wolfson Unit for Endoscopy, St Mark's Hospital and Academic Institute, Imperial College London, Harrow, London

² Department of Radiology, St Mark's Hospital and Academic Institute, Imperial College London, Harrow, London
³ Department of Haematology, Northwick Park Hospital, Harrow, London

References

- 1 *Wotherspoon AC*. Extragastic MALT lymphoma. *Gut* 2002; 51: 148–149
- 2 *Freeman C, Berg JW, Cutler SJ*. Occurrence and prognosis of extranodal lymphomas. *Cancer* 1972; 29: 252–260

- 3 *Zullo A, Hassan C, Cristofari F et al*. Effects of *Helicobacter pylori* eradication on early stage gastric mucosa-associated lymphoid tissue lymphoma. *Clin Gastroenterol Hepatol* 2010; 8: 105–110
- 4 *Isomoto H, Kamihira S et al*. A case of mucosa-associated lymphoid tissue lymphoma of the ampulla of Vater: successful treatment with radiation therapy. *Eur J Gastroenterol Hepatol* 2003; 15: 1037–1041
- 5 *Tai C-M, Tu C-H, Wu H-B et al*. An unexpected cause of chronic diarrhoea. *Gut* 2008; 57: 902, 921

Bibliography

DOI <http://dx.doi.org/10.1055/s-0031-1291744>
 Endoscopy 2012; 44: E157–E158
 © Georg Thieme Verlag KG
 Stuttgart · New York
 ISSN 0013-726X

Corresponding author

A. Murino, MD
 Wolfson Unit for Endoscopy
 St Mark's Hospital and Academic Centre
 Watford Road
 Harrow
 Middlesex
 HA1 3UJ
 UK
 Fax: +44-1702-444224
 albertomurino@yahoo.it