Rectovesical fistula treated by glue injection plus endoclippping technique

A 74-year-old man was referred to our department for endoscopic evaluation and treatment of a rectovesical fistula. He had undergone prostatectomy for early prostate cancer 2 weeks previously. Post-operatively, he had developed pneumaturia and dysuria; computed tomography (CT) had revealed a rectovesical fistula.

At colonoscopy, a well-circumscribed opening of a fistula tract was seen in the anterior rectal wall, located 2 cm beyond the linea dentata (● Fig. 1). The edges of the fistula were cauterized with argon plasma coagulation to stimulate an inflammatory reaction and local collagen synthesis. A 2-mL syringe containing 1 mL n-2-butyl cyanoacrylate (Histoacryl; B. Braun, Melsungen, Germany) and 1 mL Lipiodol was prepared and the glue/Lipiodol mix was injected via a 23-G variceal needle in aliquots of 0.5 mL at four sites of the wall of the fistula, followed by a flush of Lipiodol equivalent in volume to the dead space of the needle. The margins of the fistula orifice were brought into contact and closed with endoclips (HX-600-900; Olympus, Athens, Greece; ● Fig. 2). After endoscopic intervention, the patient was treated with parenteral nutrition and antibiotics. His symptoms completely resolved and he was discharged 7 days after the procedure. A follow-up endoscopy performed 1 month later revealed that the lesion was well healed (● Fig. 3). He remains asymptomatic 6 months post-procedure. Rectal injury after radical prostatectomy with subsequent formation of rectovesical fistula has an incidence of 1–11% [1]. York-Mason, modified York-Mason and less invasive repair methods, which include laparoscopic and robotic-assisted techniques, have been reported as successful treatment modalities for rectovesical fistulas [2,3].

We wish to emphasize the use of the combination of endoclips plus n-2-butyl cyanoacrylate injection in the treatment of gastrointestinal fistulas. To our knowledge, the present case is the first report of a rectovesical fistula that was success-

fully treated with injection of glue plus endoclippping.

Endoscopy_UCTN_Code_TTT_1AQ_2AG

Competing interests: None

P. Katsinelos¹, S. Gkagkalis¹, G. Chatzimavroudis¹, C. Zavos², J. Kountouras²

¹ Department of Endoscopy and Motility Unit, G. Gennimatas General Hospital, Thessaloniki, Greece

² Second Department of Internal Medicine, School of Medicine, Aristotle University of Thessaloniki, Ippokration Hospital, Thessaloniki, Greece

References

- 1 Harpster LE, Rommel FM, Sieber PR et al. The incidence and management of rectal injury associated with radical prostatectomy in a community urology practice. *J Urol* 1995; 154: 1435–1438
- 2 Kasraeian A, Rozet F, Cathelineau X et al. Modified York-Mason technique for repair of iatrogenic rectourinary fistula: the Montsouris experience. *J Urol* 2009; 181: 1178–1183
- 3 Sotelo R, de Andrade R, Carmona O et al. Robotic repair of rectovesical fistula resulting from open radical prostatectomy. *Urology* 2008; 72: 1344–1346

Bibliography

DOI <http://dx.doi.org/10.1055/s-0032-1308945>
Endoscopy 2013; 45: E238
© Georg Thieme Verlag KG
Stuttgart · New York
ISSN 0013-726X

Corresponding author

P. Katsinelos, MD, PhD

Department of Endoscopy and Motility Unit
G. Gennimatas General Hospital
School of Medicine, Aristotle University of Thessaloniki
Thessaloniki
Greece
Fax: +30-2310-963341
gchatzimav@yahoo.gr

Fig. 1 Endoscopic view of the orifice of a fistula located in the anterior rectal wall 2 cm from the linea dentata.

Fig. 2 View during the endoscopic procedure to close the fistula using a glue injection plus endoclippping showing complete closure of the orifice of the fistula with the use of six endoclips.

Fig. 3 View during an endoscopy performed 1 month later showing that the fistula is well healed and that some residual clips remain in place.