Complete endoscopic management of tubular esophageal duplication in a young woman

A 29-year-old woman was referred to our department for endoscopic dilation of upper esophageal stricture. Dilation was performed with Savary–Gilliard dilators allowing the passage with resistance of a standard flexible video gastroscope (EG-201FP; Fujinon, Willich, Germany). Esophagogastroduodenoscopy showed a double esophageal lumen at 18 cm from the incisors. A thick bridge of intact mucosa separated the two lumens (► Fig. 1). The passage of the endoscope through the second lumen was not possible. At 32 cm, a distal defect was also found. A barium esophagogram and high-resolution computed tomography (CT) scan confirmed esophageal tubular duplication (► Fig. 2 and ► Fig. 3).

Under general anesthesia, the standard video gastroscope was pushed down to the proximal opening of the duplication. After an easy passage of a 0.035-inch guide wire (Boston Scientific, Natick, MA, USA) in the duplicated lumen, a lengthwise incision of the intraluminal bridge was performed by using a 5.5-Fr needle-knife (microKnife XL; Boston Scientific). The incision was performed step by step, from the upper to the distal end (► Fig. 4, ► Videos 1–3). The procedure was completed with dilation of the upper esophageal stricture by using a wire-guided balloon (Boston Scientific) advanced through the endoscope and expanded up to 12 mm.

Biopsies performed along the incision showed the presence of malpighian epithelium. The patient's early post-procedural course was marked by an iatrogenic mediastinal emphysema and bilateral pneumothorax, more pronounced in the left. The placement of a left chest drain led to rapid improvement. Upper endoscopy on day 20 showed two longitudinal residual folds (► Fig. 5).

Endoscopic management of esophageal duplication was reported twice previously for the cystic form [1,2]. To our knowledge, only one case of endoscopic management of a tubular esophageal duplication has previously been reported [3]. Nevertheless, the procedure was decided upon after surgical examination through a right thoracoscopy. Our case highlights the possibility of complete endoscopic

management of tubular esophageal duplication. The post-procedure pneumothorax could have been avoided by carbon dioxide insufflation [4].

Endoscopy_UCTN_Code_TTT_1AO_2AN

Competing interests: None

Fig. 1 Upper gastrointestinal endoscopy showing two esophageal lumens separated by a septum at 18 cm from the incisors.

Fig. 2 Contrast study showing tubular duplication of the esophagus.

Fig. 3 Computed tomography (CT) scan showing evidence of tubular duplication of the esophagus.

Fig. 4 Image of interventional endoscopy showing incision of the intraluminal bridge by using a needle knife.

Fig. 5 Endoscopic image of two residual folds of the duplication.

Videos 1–3

Interventional endoscopy: a lengthwise incision was made of the intraluminal bridge by using a needle knife starting from the upper end and achieved step by step. The procedure was completed with dilation of the upper esophageal stricture.

**N. Tahri¹, L. Mnif¹, L. Chtourou¹,
M. Boudabbous¹, K. Yaïch¹, H. Fourati²,
Z. Mnif², A. Amouri¹**

¹ Department of Gastroenterology, Hedi Chaker University Hospital, Sfax, Tunisia

² Department of Radiology, Hedi Chaker University Hospital, Sfax, Tunisia

References

- 1 Will U, Meyer F, Bosseckert H. Successful endoscopic treatment of an esophageal duplication cyst. *Scand J Gastroenterol* 2005; 40: 995–999
- 2 Joyce AM, Zhang PJ, Kochman ML. Complete endoscopic resection of an esophageal duplication cyst (with video). *Gastrointest Endosc* 2006; 64: 288–289
- 3 Coumaros D, Schneider A, Tsesmeli N et al. Endoscopic management of a tubular esophageal duplication diagnosed in adolescence (with video). *Gastrointest Endosc* 2010; 71: 827–830
- 4 Uemura M, Ishii N, Itoh T et al. Effects of carbon dioxide insufflation in esophageal endoscopic submucosal dissection. *Hepato-gastroenterology* 2011; 14: 115–116

Bibliography

DOI <http://dx.doi.org/10.1055/s-0032-1309705>
Endoscopy 2012; 44: E261–E262
 © Georg Thieme Verlag KG
 Stuttgart · New York
 ISSN 0013-726X

Corresponding author

L. Mnif
 Department of Gastroenterology
 Hedi Chaker University Hospital
 Route el Ain
 3029 Sfax
 Tunisia
 Fax: +216-74-243993
 leilamnif@yahoo.fr