Meckel's ileitis versus Crohn's ileitis: diagnosis using balloon enteroscopy

Fig. 1 Enteroscopic view of the ileal lumen in an 18-year-old man who presented with episodic abdominal pain and bloody diarrhea showing an ulcerated diverticulum with some friability at the lower margin.

An 18-year-old man presented with a self-limiting episode of abdominal pain and bloody diarrhea for 5 days. During the following year, he reported similar episodes. We performed a colonoscopy that showed normal colonic mucosa with erythema and a few erosions at the ileum. Histology revealed a nonspecific inflammatory infiltrate in the lamina propria. He was treated with prednisolone with initial symptomatic relief, but was admitted to our hospital 3 weeks later, during prednisolone tapering, because of recurrent symptoms and weight loss of 3 kg. He had been an active smoker since the age of 12. Blood tests showed iron deficiency anemia (hemoglobin 9.9 g/dL) and elevated C-reactive protein (CRP; 20 mg/dL). Immunological assays were negative, including anti-*Saccharomyces cerevisiae* antibodies. Serum and fecal microbiological assays were negative.

An upper gastrointestinal endoscopy was performed, which was normal, as was the duodenal histology. We decided because of his recurrent symptoms to reappraise the colonic and ileal mucosa. We proceeded to perform a retrograde single-balloon enteroscopy, which showed an ulcerated diverticulum 80 cm from the ileocecal valve with normal distal ileal mucosa (Fig. 1). The distal margin of the diverticulum was tattooed and biopsied (Video 1). Histology showed signs of active chronic inflammation.

Fig. 2 Photograph of the specimen resected during laparoscopic enterectomy showing the ileal mucosa with circumferential mucosal folds (+) and a Meckel's diverticulum that is predominantly lined by gastric mucosa with rugal folds (*).

Fig. 3 Photomicrograph of the Meckel's diverticulum with small-intestinal mucosa on the left and heterotopic gastric body-type mucosa on the right (original magnification $\times 20$; hematoxylin and eosin [H&E] stain).

He underwent a laparoscopic enterectomy including the diverticulum 2 weeks later (Fig. 2). Pathological examination showed gastric mucosa within the diverticulum, thereby confirming the diagnosis of a Meckel's diverticulum (Fig. 3).

At follow-up after 9 months, the patient remained asymptomatic.

Meckel's ileitis is an uncommon manifestation of Meckel's diverticulum [1]. Few other cases have been reported; as in this

case, it has usually been a challenging differential diagnosis with Crohn's disease [2,3]. In our case, reassessment of the ileal mucosa with balloon enteroscopy was crucial to reaching the correct diagnosis. This

Video 1

Retrograde single-balloon enteroscopy showing a Meckel's diverticulum.

technique has been reported as a helpful tool when there is high suspicion of a Meckel's diverticulum, even when technetium pertechnetate scintigraphy is negative [4–6].

Endoscopy_UCTN_Code_CCL_1AC_2AD

Competing interests: None

M. Velosa^{1,2}, H. Cardoso^{1,3}, E. Rios⁴, G. Macedo^{1,3}

¹ Serviço de Gastrenterologia, Hospital S. João, Porto, Portugal

² Serviço de Gastrenterologia, Hospital Dr. Nélio Mendonça, Funchal, Madeira, Portugal

³ Serviço de Gastrenterologia, Faculdade de Medicina da Universidade do Porto, Porto, Portugal

⁴ Serviço de Anatomia Patológica, Hospital S. João, Porto, Portugal

References

- 1 Feldman M, Friedman LS, Brandt LJ. Sleisenger and Fordtran's Gastrointestinal and Liver Disease. 8th edn. Philadelphia: Saunders Elsevier; 2006: 2075
- 2 Levesque B, Chua H, Kane S. Meckel's ileitis: not all regional enteritis is Crohn's Disease. *Inflamm Bowel Dis* 2011; 17: E45
- 3 Honda K, Mizutani T, Higuchi N et al. A Meckel's diverticulum with an ileal ulcer detected with double-balloon enteroscopy. *Endoscopy* 2007; 39: E160
- 4 Uchiyama S, Sannomiya I, Hidaka H et al. Meckel diverticulum diagnosed by double-balloon enteroscopy and treated laparoscopically: case report and review of the literature. *Surg Laparosc Endosc Percutan Tech* 2010; 20: 278–280
- 5 Huang TY, Liu YC, Lee HS et al. Inverted Meckel's diverticulum mimicking an ulcerated pedunculated polyp: detection by single-balloon enteroscopy. *Endoscopy* 2011; 43: E244–E245
- 6 Gasbarrini A, Di Caro S, Mutignani M et al. Double-balloon enteroscopy for diagnosis of a Meckel's diverticulum in a patient with GI bleeding of obscure origin. *Gastrointest Endosc* 2005; 61: 779–781

Bibliography

DOI <http://dx.doi.org/10.1055/s-0032-1309851>
Endoscopy 2012; 44: E321–E322
 © Georg Thieme Verlag KG
 Stuttgart · New York
 ISSN 0013-726X

Corresponding author

M. Velosa, MD

Caminho dos saltos n. 32
 Vilas jardins d'agua, moradia D
 9050-219, Funchal
 Madeira Island
 Portugal
 Fax: +351-291-572538
 mo.velosa@gmail.com