Hepatic penetration by stomal ulcer: rare complication of a peptic ulcer

A 57-year-old man, who had undergone a Billroth I partial gastrectomy for a duodenal ulcer 1 year previously, was hospitalized for continual epigastric pain over the past 1 month. Laboratory testing revealed anemia (hemoglobin 9.4 g/dL; normal range 13.5–16.9) and an elevated C-reactive protein level (2.22 mg/dL; normal <0.02). Liver function tests and the serum gastrin level were within normal limits. Abdominal ultrasound revealed fluid and air bubbles in the liver (● Fig. 1 a), moving between the liver and stomach through a fistula (● Fig. 1 b). Computed tomography (CT) verified the ultrasound findings (● Fig. 2). Endoscopy revealed a large ulcer in the duodenum near the anastomosis (● Fig. 3). An endoscopic biopsy of the ulcer was negative for malignancy. The patient was diagnosed as having a stomal ulcer that had penetrated the liver, and he was subsequently treated with an H2 blocker and intravenous antibiotics. The ulcer healed after 1 month of treatment (● Fig. 4). After discharge, the patient continued to take proton pump inhibitors (PPIs), and there has been no recurrence of the ulcer during the 7-year follow-up period.

The most common site of penetration by duodenal ulcers is the pancreas (52.6%), followed by the biliary tract (18.4%), gastrohepatic omentum (10.7%), liver (6.2%), and colon (1.5%) [1]. Most cases of hepatic penetration have been diagnosed intraoperatively and/or by endoscopic biopsy [1,2]. However, in our patient, the ultrasound clearly demonstrated detailed findings sufficient for diagnosis. Most cases of ulcers penetrating the liver have been treated by surgical procedures [1,2]. In addition to the present report, there have been two other recent case reports that have demonstrated the effectiveness of medical treatments such as H2 blockers and PPIs [3,4].

Endoscopy_UCTN_Code_CCL_1AB_2AZ_3AZ

Competing interests: None

Fig. 1 Abdominal ultrasound findings in a 57-year-old man with a history of Billroth I partial gastrectomy and hospitalized for continual epigastric pain. **a** Sagittal section of the upper abdomen showing an echo-free space with air bubbles (arrowhead) in the liver. **b** Coronal section of the upper abdomen showing fluid with air bubbles (arrowhead) moving between the liver and stomach through a fistula.

Fig. 2 Contrast-enhanced computed tomography (CT) showing fluid with air in the liver and a fistula between the hepatic lesion and the stomach.

Fig. 3 Endoscopic findings. **a** An ulcer (arrow) is seen in the duodenum near the suture. **b** Close-up view of the large, deep ulcer.

Fig. 4 Repeat endoscopy on day 35 showed the healed ulcer.

A. Oka¹, Y. Amano², Y. Uchida³, K. Kagawa³, Y. Tada¹, R. Kusunoki¹, N. Fukuba¹, I. Moriyama¹, T. Yuki¹, K. Kawashima¹, S. Ishihara¹, Y. Kinoshita¹

¹ Second Department of Internal Medicine, Shimane University School of Medicine, Izumo, Shimane, Japan

² Division of Endoscopy, Kaken Hospital, Internal University of Health and Welfare, Ichikawa, Chiba, Japan

³ Department of Gastroenterology, Matsue Red Cross Hospital, Matsue, Shimane, Japan

References

- 1 Haubrich WD. Complications of peptic ulcer disease. In: Bockus HE, ed. *Gastroenterology*. Volume 1, 3rd edn. Philadelphia: WB Saunders; 1976: 720–762
- 2 Somi MH, Tarzamni MK, Farhang S et al. Liver mass due to penetration of a silent duodenal ulcer. *Arch Iran Med* 2007; 10: 242–245
- 3 Park RH, Russell RI. Liver penetration by peptic ulcer. *Am J Gastroenterol* 1988; 83: 793–795
- 4 Kayacetin E, Kayacetin S. Gastric ulcer penetrating to liver diagnosed by endoscopic biopsy. *World J Gastroenterol* 2004; 10: 1838–1840

Bibliography

DOI <http://dx.doi.org/10.1055/s-0032-1309920>
Endoscopy 2012; 44: E347–E348
 © Georg Thieme Verlag KG
 Stuttgart · New York
 ISSN 0013-726X

Corresponding author

A. Oka
 Second Department of Internal Medicine
 Shimane University Hospital
 Enya-cho
 Izumo
 Shimane 693-8501
 Japan
 Fax: +81-853-20-2187
akihiko.oka@gmail.com