Bile duct adenoma causing recurrent cholangitis: diagnosis and management with targeted Spyglass access and radiofrequency ablation

A 77-year-old man was admitted for evaluation of recurrent episodes of cholangitis. He had undergone endoscopic retrograde cholangiography (ERC) several times for biliary stone disease and for cholecystectomy, and suffered from advanced Alzheimer disease. Abdominal magnetic resonance imaging (MRI) revealed dilatation of the right posterior intrahepatic duct (● Fig. 1), with aberrant drainage directly into the common hepatic duct. However, multiple attempts to reach this segment by ERC were unsuccessful (● Fig. 2). The Spyglass system was then used for selective opacification (● Fig. 3) and optical viewing of the aberrant bile duct (Boston Scientific, Natick, Massachusetts, USA) [1,2]. A reddish nodular lesion was seen obstructing the lumen (● Fig. 4), and targeted biopsies with a 3-Fr Spybite mini-forceps (Boston Scientific) confirmed the presence of an adenoma (● Fig. 5) [3]. The stricture was dilated with a Hurricane 8-mm, 4-cm balloon (Boston Scientific) and the stones were extracted. The Spyscope was removed and two 7-Fr plastic stents placed. Given the patient's poor mental and general status, we proposed carrying out endobiliary radiofrequency ablation of the stricture. This was done 4 weeks later with a wire-guided Habib EndoHPB (Emcision, London, UK) (● Fig. 6) [4]. This case illustrates the potential impact of selective cholangioscopy with the Spyglass system on diagnosis and management of indeterminate biliary strictures.

Endoscopy_UCTN_Code_TTT_1AR_2AF

Competing interests: None

**G. Mavrogenis¹, P. H. Deprez²,
J. Wallon³, P. Warzée¹**

¹ Department of Hepato-gastroenterology,
Grand Hôpital de Charleroi, Site Notre
Dame, Charleroi, Belgium

² Department of Hepato-gastroenterology,
Cliniques Universitaires Saint-Luc,
Université Catholique de Louvain,
Brussels, Belgium

Fig. 1 Magnetic resonance imaging (MRI) showing dilatation of the right posterior hepatic duct, without visible cause of obstruction, in a 77-year-old man with history of recurrent episodes of cholangitis.

Fig. 2 Failure to opacify the aberrant right hepatic duct with classical filling techniques during endoscopic retrograde cholangiography (ERC).

Fig. 3 Targeted access to the aberrant posterior right bile duct with the Spyglass system showing tumoral obstruction.

Fig. 5 Histological section of the biliary adenoma, characterized by the regular sheets of monotonous cells, with monomorphic nuclei localized to the bottom of the cytoplasm (Papanicolaou stain; magnification $\times 25$).

³ Department of Pathology, Cliniques Universitaires Saint-Luc, Université Catholique de Louvain, Brussels, Belgium

References

- 1 Chen YK. Preclinical characterization of the Spyglass peroral cholangiopancreatography-system for direct access, visualization, and biopsy. *Gastrointest Endosc* 2007; 65: 303–311
- 2 Draganov PV, Chauhan S, Wagh MS et al. Diagnostic accuracy of conventional and cholangioscopy-guided sampling of indeterminate biliary lesions at the time of ERCP: a prospective, long-term follow-up study. *Gastrointest Endosc* 2012; 75: 347–353
- 3 Rustgi AK, Kelsey PB, Guelrud M et al. Malignant tumors of the bile ducts: diagnosis by biopsy during endoscopic cannulation. *Gastrointest Endosc* 1989; 35: 248–251
- 4 Monga A, Gupta R, Ramchandani M et al. Endoscopic radiofrequency ablation of cholangiocarcinoma: new palliative treatment modality (with videos). *Gastrointest Endosc* 2011; 74: 935–937

Bibliography

DOI <http://dx.doi.org/10.1055/s-0032-1310036>
Endoscopy 2012; 44: E290–E291
 © Georg Thieme Verlag KG
 Stuttgart · New York
 ISSN 0013-726X

Corresponding author

P. Warzée
 Grand Hôpital de Charleroi
 Site Notre Dame
 3 Grand Rue
 6000 Charleroi
 Belgium
 Fax: +327-1102-779
 pwarzee@hotmail.com

Fig. 4 Optical viewing of the aberrant bile duct with the Spyglass system showing a nodular lesion obstructing the lumen.

Fig. 6 Guide wire inserted through the biliary stenosis (left) and a Habib EndoHPB catheter placed at the level of the intraductal adenoma (right).