Rectal tonsil associated with Epstein–Barr virus

Reactive polypoid proliferation of lymphoid tissue has been described along the gastrointestinal tract [1]. Endoscopically, when a non-lymphomatous lymphoid reaction resembling tonsillar tissue is present in the rectum, the polyp is named as “rectal tonsil”. Few cases have been reported in the literature [1,2], and the etiology of this rare condition is unknown [3–5].

Here we describe the case of a 58-year-old man referred to our center with a 6-month history of fresh bleeding from the rectum. A colonoscopy was performed and a delineated round, reddish, nodular aggregate measuring 4×3×3 cm was found (► Fig. 1). A polypectomy was performed (► Fig. 2), and histological examination (► Fig. 3) revealed infiltration of mucosa, submucosa, and lamina propria of the colon by numerous lymphoid follicles. Immunohistochemical markers were positive for CD20+ (B cells), and intraepithelial lymphocytes were positive for CD3+ (T cells). Epstein–Barr virus encoded RNA in situ hybridization was positive (► Fig. 3f).

At 1- and 6-month follow-up, there was no recurrence of the lesion in the rectum. This case represents the first case of rectal tonsil associated with Epstein–Barr virus infection.

Endoscopy_UCTN_Code_CCL_1AD_2AZ

Competing interests: None

P. Grube-Pagola¹, A. Canales-Kay², A. Meixueiro-Daza³, J. M. Remes-Troche³

¹ Laboratorio de Patología Digestiva, Veracruz, México

² Colorectal Surgery Private Practice, Veracruz, México

³ Laboratorio de Fisiología Digestiva y Motilidad Gastrointestinal, Instituto de Investigaciones Médico-Biológicas, Universidad Veracruzana, Veracruz, México

Fig. 1 **a** Colonoscopy revealed a well-delineated, round, reddish, nodular aggregate measuring 4×3×3 cm, with follicles resembling the tonsils. **b** High definition colonoscopy using a digital filter (i-Scan 2, Pentax) emphasized the nodular component of the lesion.

Fig. 2 **a** Macroscopic aspect of nodular aggregate with follicles. **b** Histological view of nodular and diffuse aspect using hematoxylin and eosin (H&E) stain. **c** Macroscopic histological view using CD45 immunoreaction.

References

- 1 Farris AB, Lauwers GY, Ferry JA, Zukerberg LR. The rectal tonsil: a reactive lymphoid proliferation that may mimic lymphoma. *Am J Surg Pathol* 2008; 32: 1075–1079
- 2 O'Leary AD, Sweeney EC. Lymphoglandular complexes of the colon: structure and distribution. *Histopathology* 1986; 10: 267–283
- 3 Lloyd J, Darzi A, Teare J, Goldin RD. A solitary benign lymphoid polyp of the rectum in a 51 year old woman. *J Clin Pathol* 1997; 50: 1034–1035
- 4 Agaimy A, Mudter J, Märkl B, Chetty R. Cytomegalovirus infection presenting as isolated inflammatory polyps of the gastrointestinal tract. *Pathology* 2011; 43: 440–446
- 5 Cramer SF, Romansky S, Hulbert B et al. The rectal tonsil: a reaction to chlamydial infection? *Am J Surg Pathol* 2009; 33: 483–485

Bibliography

DOI <http://dx.doi.org/10.1055/s-0032-1310140>
Endoscopy 2012; 44: E388–E389
© Georg Thieme Verlag KG
Stuttgart · New York
ISSN 0013-726X

Corresponding author

J. M. Remes-Troche, MD
Laboratorio de Fisiología Digestiva y Motilidad Gastrointestinal
Instituto de Investigaciones Médico-Biológicas
Universidad Veracruzana
Veracruz
México
Fax: +52-229-2021231
jose.remes.troche@gmail.com

► Fig. 3 see following page.

Fig. 3 **a** Residual colonic crypts; hematoxylin and eosin (H&E) stain, $\times 200$. **b** Reactive lymphocytes and isolated immunoblasts (H&E stain, $\times 800$). **c** Binucleate reactive Reed–Sternberg-like cells (H&E stain, $\times 800$). **d** Positive CD20 immunoreaction. **e** Positive CD3 immunoreaction. **f** Epstein–Barr virus-encoded RNA was positive (in situ hybridization for Epstein–Barr virus).