Nucleophilic 5-*endo-trig* Cyclization of 3,3-Difluoroallylic Ketone Enolates: Synthesis of 5-Fluorinated 2-Alkylidene-2,3-dihydrofurans

Takeshi Fujita,^a Kotaro Sakoda,^b Masahiro Ikeda,^a Masahiro Hattori,^a Junji Ichikawa^{*a}

^a Division of Chemistry, Faculty of Pure and Applied Sciences, University of Tsukuba, Tsukuba, Ibaraki 305-8571, Japan
Fax +81(29)8534237; E-mail: junji@chem.tsukuba.ac.jp

^b Department of Chemistry, Graduate School of Science, The University of Tokyo, Hongo, Bunkyo-ku, Tokyo 113-0033, Japan

Received: 11.10.2012; Accepted after revision: 08.11.2012

Abstract: 3,3-Difluoroallylic ketones readily undergo nucleophilic 5-*endo-trig* cyclization through their metal enolates to afford 5-fluorinated 2-alkylidene-2,3-dihydrofurans. O-Cyclization exclusively occurred via intramolecular substitution of the vinylic fluorines.

Key words: cyclization, fluorine, alkenes, furans, 5-*endo-trig*, ketone enolates, vinylic substitution

gem-Difluoroalkenes (1,1-difluoro-1-alkenes) have unique reactivities toward nucleophiles, which are based on their electron-deficient and highly polarized nature. They facilitate extraordinary substitution reactions, which hardly proceed in normal alkenes.¹ Difluoroalkenes readily undergo vinylic nucleophilic substitution (S_NV) via addition to electrophilic difluoromethylene carbons and subsequent fluoride elimination. We have already reported syntheses of ring-fluorinated heterocycles by conducting the S_NV reaction of difluoroalkenes in an intramolecular fashion.² As well as sp^3 heteroatom and carbon nucleophiles,³ sp^2 nucleophiles⁴ have also participated in the 6-*endo-trig* cyclization to afford six-membered heterocycles (Scheme 1, eq 1). Furthermore, the high reactivity of 1,1-difluoro-1-alkenes has even allowed normally ‘disfavored’ 5-*endo-trig* cyclization,^{5–9} which provides scaffolds for 2-fluoro-4,5-dihydroheteroles and 2-fluorobenzoheteroles (Scheme 1, eq 2).¹⁰ Addressing the next challenge to the ‘disfavored’ process, we herein demonstrate the 5-*endo-trig* cyclization with the metal enolates of 3,3-difluoroallylic ketones, which are sp^2 atom-based ambident nucleophiles and rotationally restricted around the anionic centers (Scheme 1, eq 3). This process efficiently provides 2-alkylidene-2,3-dihydrofurans¹¹ by (i) constructing the heterocyclic ring and (ii) introducing a fluorine substituent and an alkylidene group onto the prescribed ring carbon.

The starting 3,3-difluoroallylic ketones are readily accessible through the following chemoselective difluoromethylation protocol. 1,3-Ketoaldehydes **1**, the precursors of 3,3-difluoroallylic ketones **2**, were synthesized by the acylation of either morpholine enamines **3** or metal *N*-*tert*-butyl enamides prepared by deprotonation of imines **4**, followed by hydrolysis (Scheme 2).¹² Finally, difluoroal-

6-*endo-trig* cyclization with sp^3 and sp^2 nucleophiles

5-*endo-trig* cyclization with sp^3 and sp^2 nucleophiles

5-*endo-trig* cyclization with rotationally restricted sp^2 nucleophiles

Scheme 1 Intramolecular cyclization of *gem*-difluoroalkenes

lylic ketones **2** were obtained in moderate to high yield (27–86%) via difluoromethylation of ketoaldehydes **1** by a triaminophosphonium difluoromethylide, generated in situ from dibromodifluoromethane and tris(dimethylamino)phosphine (Scheme 2).¹³ Success of the exclusively selective difluoromethylation of **1** was due to the much higher reactivity of formyl groups compared to ketone carbonyl groups.

First, we sought bases suitable for the enolate formation and the subsequent 5-*endo-trig* cyclization by using difluoroallylic ketone **2a** as a model substrate (Table 1). Lithium diisopropylamide (LDA) afforded the O-cyclization product **5a** as a single isomer, albeit in low yield, while the C-cyclization product **6a** was not detected at all (Table 1, entry 1).¹⁴ Two-fold increase in the amount of LDA (2 equiv) turned out to be effective for the cyclization (Table 1, entry 2). Also, potassium hydride (1 equiv) exclusively gave **5a** and drastically improved its yield up to 79% (Table 1, entry 3). As in the case of LDA, use of doubled amounts of potassium hydride (2 equiv) was highly effective, leading to a 91% yield of the desired dihydrofuran **5a** (Table 1, entry 4). Thus, the nucleophilic 5-*endo-trig* cy-

Scheme 2 Synthesis of 3,3-difluoroallylic ketones **2**

cyclization successfully proceeded even with rotationally restricted sp^2 nucleophiles in **2a**. This is likely due to the large polarization of the $CF_2=C$ moiety.^{10a}

Table 1 Screening of Bases Suitable for 5-*endo-trig* Cyclization of **2a**

Entry	Base (equiv)	Time	Yield of 5a (%)	Yield of 6a (%)
1	LDA (1.0)	5 h	29	— ^a
2	LDA (2.0)	4 h	42	— ^a
3	KH (1.0)	2 h	79	— ^a
4	KH (2.0)	2 h	91	— ^a

^a Not detected.

The optimized conditions obtained above for **2a** were successfully applied to the cyclizations of a variety of difluoroallylic ketones **2** (Table 2).^{15,16} Ketones **2b–g**, which are dimethylated at the allylic position, gave corresponding fluorine-containing dihydrofurans **5b–g** in good to excellent yield. Difluoroallylic benzylic ketones **2e** and **2f** gave 2-benzylidene dihydrofurans **5e** and **5f**, respectively. Reactions of difluoroallylic ketones **2h–j**, which possess a cyclohexane ring at the allylic position, constructed a spirocyclic structure in **5h–j**. The reactions of $\alpha,\alpha,\alpha',\alpha'$ -tetra-substituted ketones **2g** and **2j** were sluggish under the same conditions. However, the longer reaction time or the

Table 2 5-*endo-trig* Cyclization of 3,3-Difluoroallylic Ketones **2**

Entry	Time (h)	Product	Yield of 5 (%)
1	2	5a	91
2	2	5b	97
3	2	5c	83
4	2	5d	75
5	2	5e	98
6	2	5f	97
7	21	5g	91
8	5	5h	91
9	2	5i	94
10 ^a	3	5j	97

^a Pyridine was used as the solvent instead of THF.

use of pyridine as the solvent instead of THF improved the yields of **5g** or **5j**, respectively. Intriguingly, dihydrofuran derivatives **5a–f** and **5i** were obtained as single isomers about the *exo* double bond, judging from ^1H NMR and ^{13}C NMR studies. The configurations of **5a–f** and **5i** were assigned as *Z*-isomers by a NOESY experiment of **5b**.¹⁷ This *Z*-selectivity in the formation of dihydrofurans **5a–f** and **5i** is interpreted as follows: the *Z*-enolates seem to be generated predominantly by deprotonation of difluoroallylic ketones **2** because of steric repulsion between substituents at both of the α positions of the carbonyl groups in **2**. The subsequent cyclization presumably proceeds through the *Z*-enolates with retention of stereochemistry.

Difluoroallylic ketones **2**, as shown in Table 2, underwent 5-endo-trig O-cyclization via their enolate forms. The reaction afforded the corresponding 2-alkylidene-5-fluoro-2,3-dihydrofurans **5** without the formation of C-cyclization products, 3-fluorocyclopent-3-en-1-ones **6**. Although 5-endo-trig cyclization is assigned as disfavored in Baldwin's rules,⁵ the reactivity of 1,1-difluoro-1-alkenes allows the substrates to undergo such an extraordinary cyclization.

In summary, we have demonstrated that 3,3-difluoroallylic ketone enolates exclusively underwent intramolecular O-alkenylation to afford fluorinated dihydrofurans **5** bearing a *Z*-*exo*-alkylidene unit. The cyclization proceeded in a 5-endo-trig fashion, which is disfavored according to Baldwin's rules. In this process, a fluorine substituent was introduced selectively onto the 5-position of the 2,3-dihydrofuran scaffold. Furthermore, since fluorinated 2-alkylidene-2,3-dihydrofurans are unprecedented and highly functionalized, it is expected that these compounds would serve as parts of bioactive molecules and versatile intermediates.¹⁸

Supporting Information for this article is available online at <http://www.thieme-connect.com/ejournals/toc/synlett>.

References and Notes

- (1) For recent reviews, see: (a) Uneyama, K. *Organofluorine Chemistry*; Chap. 2.3; Blackwell Publishing: Oxford, **2006**. (b) Amii, H.; Uneyama, K. *Chem. Rev.* **2009**, *109*, 2119.
- (2) (a) Ichikawa, J. In *Fluorine-Containing Synthons, ACS Symposium Series 911*; Chap. 14; Soloshonok, V. A., Ed.; Oxford University Press/ACS: Washington DC, **2005**. (b) Ichikawa, J. *Chim. Oggi* **2007**, *25* (4), 54.
- (3) (a) Wada, Y.; Ichikawa, J.; Katsume, T.; Nohiro, T.; Okauchi, T.; Minami, T. *Bull. Chem. Soc. Jpn.* **2001**, *74*, 971. (b) Wada, Y.; Mori, T.; Ichikawa, J. *Chem. Lett.* **2003**, *32*, 1000. (c) Mori, T.; Ichikawa, J. *Chem. Lett.* **2004**, *33*, 590. (d) Ichikawa, J.; Sakoda, K.; Moriyama, H.; Wada, Y. *Synthesis* **2006**, 1590.
- (4) (a) Ichikawa, J.; Wada, Y.; Miyazaki, H.; Mori, T.; Kuroki, H. *Org. Lett.* **2003**, *5*, 1455. (b) Ichikawa, J.; Mori, T.; Miyazaki, H.; Wada, Y. *Synlett* **2004**, 1219. (c) Ichikawa, J.; Wada, Y.; Kuroki, H.; Mihara, J.; Nadano, R. *Org. Biomol. Chem.* **2007**, *5*, 3956.
- (5) For Baldwin's rules, see: (a) Baldwin, J. E. *J. Chem. Soc., Chem. Commun.* **1976**, 734. (b) Baldwin, J. E.; Cutting, J.; Dupont, W.; Kruse, L.; Silberman, L.; Thomas, R. C. *J. Chem. Soc., Chem. Commun.* **1976**, 736. (c) Baldwin, J. E.; Thomas, R. C.; Kruse, L.; Silberman, L. *J. Org. Chem.* **1977**, *42*, 3846.
- (6) Ichikawa, J.; Iwai, Y.; Nadano, R.; Mori, T.; Ikeda, M. *Chem. Asian J.* **2008**, *3*, 393; and references cited therein.
- (7) For recent reports on nucleophile-driven 5-endo-trig cyclization, see: (a) Anderson, J. C.; Davies, E. A. *Tetrahedron* **2010**, *66*, 6300. (b) Motto, J. M.; Castillo, Á.; Greer, A.; Montemayer, L. K.; Sheepwash, E. E.; Schwan, A. L. *Tetrahedron* **2011**, *67*, 1002.
- (8) For recent reports on electrophile-driven 5-endo-trig cyclization, see: (a) Stojanović, M.; Marković, R. *Synlett* **2009**, 1997. (b) Kalamkar, N. B.; Kasture, V. M.; Dhavale, D. D. *Tetrahedron Lett.* **2010**, *51*, 6745. (c) Saczewski, J.; Gdaniec, M.; Bednarski, P. J.; Makowska, A. *Tetrahedron* **2011**, *67*, 3612.
- (9) For recent reports on radical-initiated 5-endo-trig cyclization, see: (a) Pattarozzi, M.; Ghelfi, F.; Roncaglia, F.; Pagnoni, U. M.; Parsons, A. F. *Synlett* **2009**, 2172. (b) Yu, J.-D.; Ding, W.; Lian, G.-Y.; Song, K.-S.; Zhang, D.-W.; Gao, X.; Yang, D. *J. Org. Chem.* **2010**, *75*, 3232.
- (10) For 5-endo-trig cyclization of substrates with difluoroalkene moieties, see: (a) Ichikawa, J.; Wada, Y.; Fujiwara, M.; Sakoda, K. *Synthesis* **2002**, 1917. (b) Ichikawa, J.; Nadano, R.; Mori, T.; Wada, Y. *Org. Synth.* **2006**, *83*, 111. (c) Ichikawa, J. *Org. Synth.* **2011**, *88*, 162. (d) Ichikawa, J.; Wada, Y.; Okauchi, T.; Minami, T. *Chem. Commun.* **1997**, 1537. (e) Ichikawa, J.; Fujiwara, M.; Wada, Y.; Okauchi, T.; Minami, T. *Chem. Commun.* **2000**, 1887. (f) Tanabe, H.; Ichikawa, J. *Chem. Lett.* **2010**, *39*, 248. (g) Fuchibe, K.; Takahashi, M.; Ichikawa, J. *Angew. Chem. Int. Ed.* **2012**, *51*, 12059.
- (11) For selected examples of conventional synthetic methodologies for 2-alkylidene-2,3-dihydrofurans, see: (a) Tiecco, M.; Testaferri, L.; Tingoli, M.; Marini, F. *J. Org. Chem.* **1993**, *58*, 1349; and references cited therein. (b) Lattanzi, A.; Sagulo, F.; Scettri, A. *Tetrahedron: Asymmetry* **1999**, *10*, 2023. (c) Fang, Y.; Li, C. *Chem. Commun.* **2005**, 3574. (d) Chen, Y.-F.; Wang, H.-F.; Wang, Y.; Luo, Y.-C.; Zhu, H.-L.; Xu, P.-F. *Adv. Synth. Catal.* **2010**, *352*, 1163. (e) Montel, S.; Bouyssi, D.; Balme, G. *Adv. Synth. Catal.* **2010**, *352*, 2315.
- (12) For the synthesis of 1,3-ketoaldehydes from enamines, see: Kuhlmei, S.-R.; Adolph, H.; Rieth, K.; Opitz, G. *Liebigs Ann. Chem.* **1979**, 617.
- (13) For difluoromethylation of aldehydes with dibromodifluoromethane and tris(trimethylamino)phosphine, see: (a) Nae, D. G.; Burton, D. J. *Synth. Commun.* **1973**, *3*, 197. (b) Vinson, W. A.; Prickett, K. S.; Spahic, B.; deMontellano, P. R. O. *J. Org. Chem.* **1983**, *48*, 4661.
- (14) Baldwin noted that when endocyclic alkylation of ketone enolates constructs five-membered rings, O-cyclization would be preferable because of an in-plane approach to enolates. The chemoselectivity in our case could be partially explained by a similar reasoning, albeit with the $\text{sp}^2\text{-CF}_2$ electrophile instead of $\text{sp}^3\text{-C}$ electrophiles. See: Baldwin, J. E.; Kruse, L. I. *J. Chem. Soc., Chem. Commun.* **1977**, 233.
- (15) **(Z)-5-Fluoro-3,3-dimethyl-2-(2-phenylethylidene)-2,3-dihydrofuran (5b)**
To a suspension of KH (oil free, 46 mg, 1.2 mmol) in THF (11 mL) was added 6,6-difluoro-4,4-dimethyl-1-phenylhex-5-en-3-one (**2b**, 138 mg, 0.58 mmol), and the mixture was heated to reflux for 2 h. After cooling to r.t., the reaction was quenched with phosphate buffer (pH 7). Organic materials were extracted with Et_2O three times. The combined extracts

were washed with brine and dried over MgSO_4 . After removal of the solvent under reduced pressure, the residue was purified by TLC on silica gel (EtOAc–hexane, 1:5) to give **5b** (122 mg, 97%) as a colorless oil. ^1H NMR (500 MHz, CDCl_3): δ = 1.26 (d, J_{HF} = 1.1 Hz, 6 H), 3.45 (d, J = 7.5 Hz, 2 H), 4.20 (d, J_{HF} = 5.4 Hz, 1 H), 4.73 (td, J = 7.5 Hz, J_{HF} = 3.4 Hz, 1 H), 7.18–7.22 (m, 3 H), 7.27–7.30 (m, 2 H). ^{13}C NMR (126 MHz, CDCl_3): δ = 30.1 (d, J_{CF} = 2 Hz), 30.9, 44.2 (d, J_{CF} = 2 Hz), 79.7 (d, J_{CF} = 8 Hz), 99.4, 125.9, 128.2, 128.4, 141.0, 157.5 (d, J_{CF} = 276 Hz), 160.6 (d, J_{CF} = 3 Hz). ^{19}F NMR (470 MHz, CDCl_3): δ = 46.2 (s). IR (neat): 3028, 2970, 2931, 1801, 1726, 1703, 1454, 1279, 1219, 1126, 1088, 993, 976, 748, 698 cm^{-1} . Anal. Calcd for $\text{C}_{14}\text{H}_{15}\text{FO}$: C, 77.04; H, 6.93. Found: C, 76.80; H, 7.16%.

- (16) 3,3-Disubstituted 5-fluoro-2-alkylidene-2,3-dihydrofurans **5** are air- and heat-stable.

- (17) In the NOESY experiment of dihydrofuran **5b**, substantial correlation between the methyl protons and the vinylic proton H^a was observed. No NOE correlation was detected between the methyl protons and the allylic protons H^b (Figure 1).

Figure 1 NOE correlation between protons in dihydrofuran **5b**

- (18) For a review on bioactivities of fluorinated compounds, see: Müller, K.; Faeh, C.; Diederich, F. *Science* **2007**, 317, 1881.