Spontaneous intraductal stent migration after endoscopic ultrasound-guided choledochogastrostomy

Fig. 1 Abdominal radiograph showing a partially covered self-expanding metal stent in position between the extrahepatic bile duct and the gastric antrum.

Fig. 2 Endoscopic view 1 month after formation of the choledochogastrostomy showing the partially covered self-expanding metal stent that had migrated into the bile duct.

Endoscopic ultrasound-guided biliary drainage (EUS-BD) is increasingly being reported as an alternative method of biliary decompression [1]. EUS-guided choledochogastrostomy (EUS-CGS) is considered a possible salvage therapy for patients in whom EUS-guided choledochoduodenostomy (EUS-CDS) or hepaticogastrostomy (EUS-HGS) is not possible [2]. We report a case of intraductal migration of a partially-covered self-expandable metallic stent (PCSEMS) after EUS-CGS, which was managed successfully with endoscopic intervention.

A 56-year-old woman with locally advanced cancer of the pancreatic head was admitted to our hospital. She had previously undergone a gastrojejunostomy and percutaneous biliary drainage for malignant gastric outlet obstruction and biliary obstruction at another hospital. Given her strong desire to have the exter-

nal drainage tube removed and because EUS-CDS and EUS-HGS were impossible for anatomical reasons, she underwent EUS-CGS with insertion of a PCSEMS for internal biliary drainage without complications (● Fig. 1; ● Video 1).

The external drainage tube was removed successfully 15 days after EUS-CGS, but 1 month later she developed acute cholangitis. An emergency endoscopy revealed that the PCSEMS had migrated into the bile duct (● Fig. 2), which was confirmed on an abdominal radiograph (● Fig. 3). The remaining fistula was successfully cannulated and a guidewire was advanced through the migrated PCSEMS into the intrahepatic bile duct. Another PCSEMS was placed across the fistula between the first PCSEMS and the stomach (● Fig. 4; ● Video 2). The cholangitis subsided and she was discharged 3 days after the procedure without complications.

Spontaneous intraductal migration of a covered metallic stent after EUS-BD is a serious complication [3]. We managed this successfully with endoscopic placement of a further PCSEMS across the resulting fistula. Because the extrahepatic bile duct and gastric antrum are not connected anatomically, we must be aware of this complication after EUS-CGS.

Endoscopy_UCTN_Code_CPL_1AK_2AJ

Competing interests: None

Video 1

Endoscopic ultrasound-guided choledochogastrostomy (EUS-CGS) using a partially-covered self-expandable metallic stent (PCSEMS).

Video 2

Placement of a second partially-covered self-expandable metallic stent (PCSEMS), which was positioned across the fistula between the first PCSEMS and the stomach.

Fig. 3 Abdominal radiograph showing the distal end of the partially-covered self-expandable metallic stent located inside the fistula.

Fig. 4 Abdominal radiograph showing a second partially-covered self-expandable metallic stent (PCSEMS) placed across the fistula between the first PCSEMS and the stomach.

**K. Kawakubo, H. Kawakami,
M. Kuwatani, S. Haba, T. Kudo,
Y. Abe, N. Sakamoto**

Department of Gastroenterology
and Hepatology, Hokkaido University
Graduate School of Medicine, Sapporo,
Japan

References

- 1 Isayama H, Nakai Y, Kawakubo K et al. Recent progress in endoscopic ultrasonography guided biliary intervention. *Clin J Gastroenterol* 2012; 2: 93–100
- 2 Artifon EL, Okawa L, Takada J et al. EUS-guided choledochostomy: an alternative for biliary drainage in unresectable pancreatic cancer with duodenal invasion. *Gastrointest Endosc* 2011; 73: 1317–1320
- 3 Martins FP, Rossini LG, Ferrari AP. Migration of a covered metallic stent following endoscopic ultrasound-guided hepaticogastrostomy: fatal complication. *Endoscopy* 2010; 42: E126–E127

Bibliography

DOI <http://dx.doi.org/10.1055/s-0032-1326255>
Endoscopy 2013; 45: E89–E90
© Georg Thieme Verlag KG
Stuttgart · New York
ISSN 0013-726X

Corresponding author

H. Kawakami, MD
Department of Gastroenterology and Hepatology
Hokkaido University Graduate School of Medicine
Kita 15, Nishi 7, Kita-ku
Sapporo 060-8638
Japan
Fax: +81-11-7067867
hiropon@med.hokudai.ac.jp