Gastric balloon causing small bowel obstruction: treatment by double-balloon enteroscopy

A 32-year-old woman was referred to our emergency department because of abdominal pain and vomiting. She had received a gastric balloon in an outpatient clinic 5 years previously to treat obesity, but had not attended follow-up appointments. Emergency computed tomography showed a partially deflated gastric balloon (● Fig. 1, arrow) in the mid part of the jejunum with proximal dilatation of the small bowel loops consistent with ileus.

To remove the gastric balloon from the jejunum, the patient was offered peroral double-balloon enteroscopy under conscious sedation. The gastric balloon was found at an insertion depth of 180 cm (● Fig. 2a). To empty the methylene-blue-containing saline out of the gastric balloon, a standard injection needle was used to

perforate the silicone wall (● Fig. 2b). The deflated gastric balloon was folded up using a large polypectomy snare and was removed (● Fig. 2c). The symptoms resolved quickly and the patient was discharged the next day.

Insertion of gastric balloons is performed as a temporary measure for weight loss [1]. If a gastric balloon remains longer than scheduled, it may deflate spontaneously and cause obstruction by entering the bowel. Similar cases have been treated by surgery [2–5]. However, double-balloon enteroscopy seems a promising method to treat bowel obstruction caused by partially deflated gastric balloons.

Endoscopy_UCTN_Code_CPL_1AH_2AK

Competing interests: None

Fig. 1 CT scan of the abdomen of a 32-year-old woman showing dilated small-bowel loops and the partially deflated gastric balloon (arrow).

Fig. 2 a–c Removal of the gastric balloon by double-balloon enteroscopy: **a** gastric balloon in the mid part of the jejunum accompanied by superficial ulcerations; **b** perforation of the silicone wall with an injection needle; **c** removal of the gastric balloon with a polypectomy snare.

U. Halm, M. Grothoff, R. Lamberts

Park-Krankenhaus Leipzig, Klinik für Innere Medizin II, Leipzig, Germany

References

- 1 *Dumonceau JM*. Evidence-based review of the Bioenterics intragastric balloon for weight loss. *Obes Surg* 2008; 18: 1611–1617
- 2 *Kim WY, Kirkpatrick UJ, Moody AP et al*. Large bowel impaction by the BioEnterics intragastric balloon (BIB) necessitating surgical intervention. *Ann R Coll Surg Engl* 2000; 82: 202–204
- 3 *Henninger C, Kramer M, Horger M et al*. Ileus of uncommon cause – important and weighty facts. *Dtsch Med Wochenschr* 2009; 134: 2640
- 4 *Hegade VS, Sood R, Douds AC*. Small bowel obstruction induced by spontaneous partial deflation of an intragastric balloon. *Ann R Coll Surg Engl* 2012; 94: e171–e173
- 5 *Moszkowicz D, Lefevre JH*. Deflated intragastric balloon-induced small bowel obstruction. *Clin Res Hepatol Gastroenterol* 2012; 36: e17–e19

Bibliography

DOI <http://dx.doi.org/10.1055/s-0032-1326265>
Endoscopy 2013; 45: E78–E79
 © Georg Thieme Verlag KG
 Stuttgart · New York
 ISSN 0013-726X

Corresponding author**U. Halm, MD**

Park-Krankenhaus Leipzig
 Strümpellstr. 41
 04289 Leipzig
 Germany
 Fax: +49-341-8642264
ulrich.halm@parkkrankenhaus-leipzig.de