Early neoplasia of the common bile duct diagnosed and completely removed using multiple endoscopic modalities

Fig. 1 Endoscopic ultrasound (EUS) view showing a 5-mm polypoid mass in the mid to distal portion of the common bile duct.

Fig. 2 Endoscopic view during endoscopic retrograde cholangiopancreatography (ERCP) showing the polypoid mass that popped out during balloon extraction.

Fig. 3 Histopathology of the removed polypoid mass showing high grade dysplasia and a glandular pattern without definite stromal invasion.

We report the case of a 73-year-old man who presented with a recurrent liver abscess, the second episode occurring after an interval a few months. Blood cultures were positive for *Klebsiella pneumoniae* on both admissions. He underwent magnetic resonance cholangiopancreatography (MRCP), which revealed a mildly dilated common bile duct (CBD) without any definite cause of obstruction. He was therefore scheduled to undergo further evaluation by endoscopic ultrasound (EUS), for which he was sedated in the endoscopic suite using intravenous propofol. Linear EUS (EG-3870UTK, Pentax, Miyaki, Japan) showed a 5-mm polypoid mass in the mid to distal portion of the CBD (► Fig. 1). The gallbladder showed

evidence of chronic cholecystitis without evidence of gallstones.

Endoscopic retrograde cholangiopancreatography (ERCP) was performed next and revealed a mildly dilated CBD with no definite filling defect. The CBD was swept using an extraction balloon (B-V233P-A/B, Olympus Medical System Co., Tokyo, Japan), and during this procedure, a pinkish polypoid 5-mm mass popped out (► Fig. 2). This lesion was removed using biopsy forceps and histopathology showed it to be an adenoma with high grade dysplasia and an ill-defined margin (► Fig. 3a). As a result, the patient was scheduled for a cholangioscopy to further evaluate the base of the adenoma.

Direct per oral cholangioscopy was performed using an ultraslim gastroscope (GIF-N180, Olympus), and the residual stalk of the polyp was identified in the mid to distal CBD (► Fig. 4a, b). Narrow band imaging showed normal mucosa and a normal vascular pattern at the base of the lesion (► Fig. 4c, d). Multiple biopsies were however obtained, histopathology of which showed normal bile duct epithelium without evidence of residual neoplasm.

CBD adenomas are rare benign neoplasms of the bile duct [1–3], although there has been a report of a bile duct adenoma giving rise to an adenocarcinoma [4]. To date, there have been only a few case reports on the endoscopic treatment of bile duct adenomas. In our patient, the lesion was a premalignant CBD adenoma that was successfully treated using multiple endoscopic modalities.

Endoscopy_UCTN_Code_TTT_1AR_2AF

Competing interests: None

**N. Sirimontaporn¹, P. Aswakul¹,
P. Junyangdikul², V. Prachayakul³**

¹ Liver and Digestive Institute, Samitivej Sukhumvit Hospital, Bangkok, Thailand

² Department of Pathology, Samitivej Srinakarin Hospital, Bangkok, Thailand

³ Siriraj GI Endoscopy Center, Siriraj Hospital, Bangkok, Thailand

Fig. 4 Images obtained during cholangioscopy to further evaluate the stalk of the polyp in the common bile duct showing: **a** the initial cholangioscopic view; **b** a fluoroscopic image of the ultraslim gastroscopy in the common bile duct **c** a cholangioscopic view of the biopsy site; **d** a cholangioscopic view of the biopsy site with narrow band imaging, which revealed normal mucosa and vascular pattern.

References

- 1 Yang J, Wang W, Yan L. The clinicopathological features of intraductal papillary neoplasms of the bile duct in a Chinese population. *Dig Liv Dis* 2012; 44: 251–256
- 2 Koga F, Tanaka H, Takamutsu S et al. A case of very large intrahepatic bile duct adenoma followed for 7 years. *World J Clin Oncol* 2012; 3: 63–66
- 3 Saleem A, Wang KK, Baron TH. Successful endoscopic treatment of intraductal extension of a villous adenoma with high-grade dysplasia, with 3-years follow-up. *Gastro-intest Endosc* 2011; 74: 714–716
- 4 Munshi AG, Hassan M. Common bile duct adenoma: case report and brief review of literature. *Surg Laparosc Endosc Percutan Tech* 2010; 20: e193–e194

Bibliography

DOI <http://dx.doi.org/10.1055/s-0032-1326278>
Endoscopy 2013; 45: E102–E103
 © Georg Thieme Verlag KG
 Stuttgart · New York
 ISSN 0013-726X

Corresponding author

V. Prachayakul, MD
 Division of Gastroenterology,
 Department of Internal Medicine
 Siriraj Hospital
 Mahidol University
 Bangkok 10700
 Thailand
 Fax: +66-2419-6101
kaiyjr@gmail.com