Acute suppurative pancreatic ductitis associated with pancreatic duct obstruction

Fig. 1 Abdominal plain computed tomography (CT) in a 70-year-old man with abdominal pain and high fever, showing a 10-mm calcified stone in the head of the pancreas.

Fig. 2 Endoscopic view showing discharge of a large amount of pus mixed with pancreatic juice after a catheter was inserted in the papilla of Vater.

A 70-year-old man with abdominal pain and high fever was referred to our hospital because of suspected branch duct intraductal papillary mucinous neoplasm and pancreatolithiasis. Laboratory results showed a total bilirubin of 4.7 mg/dL, slight elevation in the serum pancreatic enzyme level and marked elevation in the serum C-reactive protein level and white blood cell count. Plain computed tomography (CT) showed mild swelling of the pancreas and a 10-mm calcified stone in the head of the pancreas (● Fig. 1). Due to the presence of obstructive cholangitis and pancreatitis, endoscopic retrograde cholangiopancreatography was carried out for drainage. When a catheter was inserted in the papilla of Vater, there was discharge of a large amount of pus mixed

Fig. 3 A cannula was inserted deep into the common bile duct and the pancreatic duct, and bile and pancreatic juice were aspirated, respectively. The pancreatic juice was purulent (left tube), but the bile was normal (right tube).

with pancreatic juice (● Fig. 2). The pancreatic juice was purulent (● Fig. 3, left tube), but the drained bile was not (● Fig. 3, right tube). *Klebsiella oxytoca* was isolated from the pancreatic juice. Following nasobiliary drainage and medi-

cal treatment with antibiotics and gabexate mesilate, the patient recovered quickly with no fever and the abdominal pain. The final diagnosis was acute suppurative pancreatic ductitis (ASPD).

Weinman [1] reported the first case of ASPD characterized by purulent discharge. Acute suppurative cholangitis due to biliary ductal obstruction is a common disease, but there have been only a few reported cases of ASPD [2,3]. Drainage of the pancreatic duct is quite effective for relief of ASPD. Therefore an endoscopist should keep in mind the possibility of ASPD in cases suspected of having suppurative cholangitis.

Endoscopy_UCTN_Code_CCL_1AZ_2AM

Competing interests: None

T. Fujinaga, T. Nishida, M. Miyazaki, M. Shigekawa, K. Ikezawa, K. Iwahashi, T. Inoue, T. Yamada, H. Ezaki, S. Shinzaki, T. Yakushijin, H. Iijima, M. Tsujii, T. Takehara

Department of Gastroenterology and Hepatology, Osaka University Graduate School of Medicine, Osaka, Japan

References

- 1 Weinman DS. Acute suppurative of the pancreatic duct. *Gastrointest Endosc* 1995; 41: 268–270
- 2 Tajima Y, Kuroki T, Susumu S et al. Acute suppurative of the pancreatic duct associated with pancreatic ductal obstruction due to pancreas carcinoma. *Pancreas* 2006; 33: 195–197
- 3 Fujimori N, Igarashi H, Ito T. Acute obstructive suppurative pancreatic ductitis. *Clin Gastroenterol Hepatol* 2011; 9: A28

Bibliography

DOI <http://dx.doi.org/10.1055/s-0032-1326450>
Endoscopy 2013; 45: E135
© Georg Thieme Verlag KG
Stuttgart · New York
ISSN 0013-726X

Corresponding author

T. Takehara, MD, PhD
Department of Gastroenterology and Hepatology
Clinical Research Building
Osaka University Graduate School of Medicine
2-2 Yamadaoka, Suita
Osaka 565-0871
Japan
Fax: +81-6-6879-3629
takehara@gh.med.osaka-u.ac.jp